

Review Article

A contemporary therapeutic approach to bone disease in beta-thalassemia - a review

Dimitrios Stefanopoulos¹, Nikolaos A. Papaioannou¹, Athanassios G. Papavassiliou², George Mastorakos³, Andromachi Vryonidou⁴, Aikaterini Michou⁵, Ismene A. Dontas¹, George Lyritis⁶, Eva Kassi^{2,7*}, Symeon Tournis^{1*}

¹Laboratory for Research of the Musculoskeletal System "Th. Garofalidis", KAT Hospital, National and Kapodistrian University of Athens, School of Medicine, Athens, Greece; ²Department of Biological Chemistry, Medical School, National and Kapodistrian University of Athens, Athens, Greece; ³Second Department of Obstetrics and Gynecology, National and Kapodistrian University of Athens, Aretaieion Hospital, Athens, Greece; ⁴Department of Endocrinology, Diabetes & Metabolism, General Hospital Korgialenio-Benakio, Athens, Greece; ⁵Department of Endocrinology, "Elena Venizelou" General Hospital, Athens, Greece; ⁶Hellenic Osteoporosis Foundation, Athens, Greece; ⁷First Department of Internal Medicine - Medical School- Laikon Hospital - National and Kapodistrian University of Athens, Greece

* Equally contribution

Abstract

Homozygous beta-thalassemia represents a serious hemoglobinopathy, in which an amazing prolongation in the survival rate of patients has been achieved over recent decades. A result of this otherwise positive evolution is the fact that bone problems have become a major issue in this group of patients. Through an in-depth review of the related literature, the purpose of this study is to present and comment on the totality of the data that have been published to date pertaining to the prevention and treatment of thalassemia bone-disease, focusing on: the contribution of diet and lifestyle, the treatment of hematologic disease and its complications, the management of hypercalciuria, the role of vitamins and minerals and the implementation of anti-osteoporosis medical regimen. In order to comprehensively gather the above information, we mainly reviewed the international literature through the PubMed database, searching for the preventive and therapeutic data that have been published pertaining to thalassemia bone-disease over the last twenty-nine years. There is no doubt that thalassemia bone-disease is a complication of a multi-factorial etiopathology, which does not follow the rules of classical postmenopausal osteoporosis. Bisphosphonates have been the first line of treatment for many years now, with varied and usually satisfactory results. In addition, over the last few years, more data have arisen for the use of denosumab, teriparatide, and other molecules that are in the clinical trial phase, in beta-thalassemia.

Keywords: Thalassemia, Osteoporosis, Bisphosphonates, Denosumab, Activin-A

Introduction

Beta-Thalassemia (β Th) is a congenital disease in which the genes related to the production of beta-polypeptide chains of hemoglobin are damaged. It manifests as severe hemoglobinopathy with a variety of genotype forms and the degree of gene damage dictates the final ability of the body to produce from practically none at all ("Thalassemia Major") to relatively satisfactory ("Thalassemia Intermedia") beta-chain levels and normal hemoglobin HbA¹. The total population of patients with β Th is estimated today to amount to 100,000

The authors have no conflict of interest.

Corresponding author: Dimitrios Stefanopoulos, Laboratory for Research of the Musculoskeletal System "Th. Garofalidis", KAT Hospital, National and Kapodistrian University of Athens, School of Medicine, Athens, Greece Ermou 50, Nea Makri, 19005, Attica, Greece

E-mail: dsteph@endo.gr

Edited by: George Lyritis

Accepted 27 January 2018

persons worldwide^{2,3}. Clinically, the disease is characterized by ineffective erythropoiesis, extensive hemolysis, severe anemia and hepatosplenomegaly. Therapeutically, frequent blood transfusions of packed red cells are usually required, a condition which leads to especially harmful deposition of cytotoxic iron in the tissues and organs of the patients^{1,3}. As a result, regular chelation therapy is strongly recommended. The serious cardiovascular manifestations, endocrine gland disorders, the severe bone disease, the high incidence of hepatitis B and C, as well as the accompanying chronic liver disease are mentioned as only some of the numerous complications present in β Th^{1,3}.

Because of the impressive increase in patient survival rates, the bone disease which accompanies beta-thalassemia (TBD) affects the large majority of the cases in our days. It is characteristically mentioned that just the frequency of decreased bone mineral density (BMD) amounts to 40-50% of adult β Th-patients worldwide, while surpassing even 90% in older and/or inadequately treated individuals⁴⁻⁶. The spine seems to be the most affected location, followed by the distal radius and femur⁷. Separate studies have demonstrated an incidence of fractures ranging from 12% to 71%, depending on the characteristics of the population being studied^{4,5,8-10}. It has been described that these are observed mostly in the upper limbs (33-53.3%), followed by spinal, pelvic and hip fractures, with significantly lower frequency^{8,11}. Of special interest is the observation that bone complications are observed with the same frequency between the sexes, while more severe bone disease seems to develop in patients with β Th-intermedia who had been transfused at extremely sparse intervals in the past¹²⁻¹⁵. Finally, we must emphasize that TBD is clearly differentiated from classical postmenopausal osteoporosis. Although low BMD and typical osteoporotic changes in the bone architecture are present, these changes develop at much younger ages -often as early as the second decade of life- while additional disorders also coexist¹⁶. Indicatively, mention is made of abnormal growth of the long bones, hyperplasia of the erythropoietic bone marrow, osteomalacia and metabolic bone disorders arising from the endocrine complications of the disease^{7,17,21}.

Methods

In order to carry out this research, we reviewed the international literature from 1988 to 2017 via the PubMed database. As keywords, we used the terms: Thalassemia AND Bone disease (or Thalassemia AND Osteoporosis) AND Management, Treatment, Hypercalciuria, Calcium, Vitamins, Zinc, Bisphosphonates, Denosumab, Teriparatide, Activin-A. We focused on twenty-five interventional trials and nine older reviews pertaining to the pathogenesis, prevention and treatment of TBD. We also consulted medical books specific to β Th, with significant references to the skeletal complications of the disease. Finally, we utilized information on the results to date of phase-III interventional clinical trials that are underway, as these were presented at 21st

and 22nd annual congresses of the European Hematology Association (EHA).

Causes – Pathophysiology

TBD exhibits an extremely multifactorial etiology, something that complicates both the understanding of the development of the disorder, as well as its diagnosis and treatment. The purpose of this review is not to provide a detailed list and thorough description of those pathogenic pathways. Therefore, the etiology and pathogenesis of the bone disease will be described briefly, in order for the therapeutic approaches being implemented today to be better understood.

a) Hyperplasia of the erythropoietic bone marrow. The conditions of chronic hypoxia that prevail in the bodies of β Th-patients lead to an excessive expansion in the erythropoietic bone marrow. In the past, this mechanism resulted in marked deformation of bones rich in erythropoietic marrow, such as the cancellous bones of the skull, the vertebrae, as well as the bones of the pelvis⁵. Today, the therapeutic protocols of frequent blood transfusions suppress the marrow sufficiently. However, a degree of increased activity continues to exist, leading to degeneration and thinning of the trabeculae, thinning of the cortex and, eventually, alterations of the normal bone architecture with degeneration of the bone quality^{5,15,22,23}.

b) Genetic predisposition. Genetic disorders that are diagnosed with increased frequency in the context of β Th have been considered responsible for modification of the genetic expression in a manner that favors the development of TBD in these patients. Polymorphisms of COLIA1 and COLIA2 genes (which affect the quality of collagen), polymorphisms such as BsmI and FokI (which affect the function of the vitamin-D receptor), polymorphisms affecting estrogen-receptors ER α and ER β and calcitonin-receptor CALC-R, and even disorders in the sequencing of TGF- β 1, have been implicated^{4,5,7,18}.

c) Reduced physical activity. Often, β Th-patients do not exercise sufficiently, either due to serious co-morbidities or even due to overprotective parenting at younger ages^{14,19}.

d) Hemosiderosis. Excessive iron deposition in the bone tissue: i) exerts a direct toxic effect on osteoblasts reducing their number, ii) stimulates the secretion of Receptor Activator of Nuclear factor-kappaB Ligand (RANKL) and iii) inhibits the physiological mineralization of the osteoid^{5,24-28}. Furthermore, hemosiderosis exerts additional harmful effects on the skeleton through indirect mechanisms, affecting the functioning of other tissues and organs as discussed in detail below.

e) Toxicity due to chelating agents. Trials exist which prove the positive impact of chelating agents on bone metabolism due to a reduction in the iron overload⁶⁹. However, when these elements are improperly utilized, administered in high doses and/or without proper evaluation of the patients, this may lead to the development of several negative effects, such as pathological growth of the long

bones, platyspondyly, arthropathy, a disruption in normal functioning of the OPG/RANK/RANKL system in favor of osteoclasts, a reduction in the proliferation of osteoblasts and fibroblasts, as well as a lack in bone protective minerals and trace elements like calcium, phosphorus, zinc and vitamin-D^{2,5,24,29-34}.

f) The OPG/RANK/RANKL system. A disruption in the normal balance of the ratio of the soluble fraction of RANKL to osteoprotegerin (OPG) has been observed in β Th, resulting in an increase in the sRANKL/OPG ratio and bone resorption. This has mainly to do with an overproduction of RANKL and, secondarily, with a possible slight reduction in OPG levels. In term of etiopathogenesis, hemosiderosis, chelating agents, hypogonadism and neurosecretory dysfunction of the growth hormone (GH) / insulin-like growth factor-1 (IGF-1) axis are implicated^{4,17,35}.

g) Other cytokines and growth factors. Elevated levels of cytokines and growth factors that promote bone resorption are often detected in the sera of patients with β Th. Those molecules which present the greatest increase and have clearly been implicated in inducing bone complications in β Th are interleukins 1 and 6, sclerostin and the factors Dickkopf-1 and TNF- α ^{5,36,37}.

h) Endocrine and metabolic disorders. The endocrine glands are organs with extremely heightened sensitivity to the cytotoxic effects of iron overload. This fact leads many β Th-patients to develop a plethora of endocrine complications, which further affect their already disturbed bone metabolism.

- (i) *Hypogonadism* is due to a malfunction of the gonadotrophic cells of the pituitary gland and/or the cells of the gonads (testes-ovaries). In related trials, it is reported that the complication is observed in 30-55% of patients of both sexes and often begins at a relatively young age^{4,5,38-42}.
- (ii) *GH/IGF-1 axis disorder.* This is due to a disorder of the somatotrophic cells of the pituitary gland as well as possible concomitant hepatopathy. The incidence of the complication has not been firmly established, as most trials indicate that it usually develops in 7.9-15% of all patients, although higher rates of up to 32% have also been reported⁴³⁻⁴⁶.
- (iii) *Diabetes mellitus* in β Th usually begins as a kind of type-2 diabetes in the second decade of life or later. This occurs at a rate of 7-11%, while prediabetes is seen with much greater frequency^{47,48}.
- (iv) *Hypothyroidism.* Thyroid function disorders occur in approximately 10-15% of β Th-patients and are related almost entirely to the presence of subclinical or clinical hypothyroidism due to damage to the cells of the thyroid and/or the thyrotroph cells of the pituitary gland^{4,5,49}.
- (v) *Hypoparathyroidism* affects 13.5-14.6% of patients and can lead to severe hypocalcemia, hypercalciuria and inadequate production of active vitamin-D^{50,51}.
- (vi) *Vitamin deficiencies.* Beta-thalassemia often co-

exists with liver dysfunction, impaired kidney function, and malabsorption due to hemosiderosis of the gastrointestinal tract. Combined with the effects of chelating agents, the above conditions have been deemed responsible for the very common vitamins D and C deficiencies seen in β Th-patients. Indicatively, values of 25OHD <20 ng/ml have been described at a rate of greater than 42%, and between 20-29 ng/ml in 30% of American patients, mainly of Asian and Caucasian descent^{5,24,29,52-54}.

- (vii) *Metal homeostasis disturbance:* In approximately 30% of adult patients hypercalciuria develops, often complicated by nephrolithiasis, which is estimated to be present in up to 18% of patients²³. Indeed, these percentages seem to increase more when the population being studied pertains to adult patients treated with the chelating agent deferasirox in moderate and high doses. The pathogenesis of the disorder is multi-factorial (chronic hypoxia, frequent blood transfusions, iron toxicity in the kidneys, chelation, hypoparathyroidism etc)^{31,32,55-58}. Hyperphosphaturia has also been reported in β Th, but at lower rates. It has been associated with chelation regimens, while it is also feasible that hemosiderosis itself may be a predisposing factor in its development, since the intravenous administration of iron seems to promote the development of phosphaturia by increasing fibroblast growth factor-23 FGF-23 in the serum^{16,59,60}. In addition to the above, albuminuria and hyperuricosuria are also observed more often in β Th than in the general population. Hypocalcemia represents yet another manifestation that seems to develop more frequently in the patients and is due mainly to hypoparathyroidism. Finally, low levels of other minerals have been observed in the blood, such as magnesium, copper and zinc, a condition which is attributed to both the action of chelation as well as malabsorption in the gastrointestinal tract^{5,29,61}.

A contemporary therapeutic approach

As the development of TBD is attributed to numerous causes and disorders, methodical and comprehensive treatment should ideally be directed towards all the factors that affect the skeleton in β Th. Nowadays, it has become clear that any treatment intervention must begin promptly and continually be adjusted depending on changes in the clinical history of each patient over time. The therapeutic approach to TBD includes the following levels of preventive and curative measures, which should ideally be combined.

A. General measures and Lifestyle

In β Th, the degree of maintenance -or even improvement- in bone mass is remarkable when patients adopt a healthy lifestyle and follow appropriate instructions as regards exercise, nutrition, and behaviors which improve bone metabolism. These general measures include^{5,61-63}: (1) The daily adequate intake of dairy products in quantities that

fall within the guidelines aimed at the normal population. A mild exposure of a portion of the skin to the sun for approximately 20 minutes daily is also recommended. (2) Restricting the excessive consumption of salt, caffeine, protein and carbonated soft-drinks which are known, in large quantities, to increase calcium excretion in the urine. (3) Restricting alcohol consumption and smoking cessation. In addition to the other complications that these habits can cause, they negatively affect bone health by disrupting the normal functioning of bone cells and the metabolism of calcium^{53,64-67}. (4) Adopting an individualized exercise regimen -in consultation with the hematologist and the cardiologist- in order to promote bone formation. Ideally, the exercise must be carried out for at least 30 minutes daily, five days per week. The benefits of exercise in TBD are unquestionable since, in addition to the direct impact on bone metabolism, exercise improves cardiovascular functioning as well as the endocrinological and metabolic profile of patients, which in turn affect bone health^{4,5,17}.

B. Treatment of hematological disease and its complications

- I. Programs of regular blood transfusions for the purpose of maintaining hemoglobin levels greater than 9-10.5 g/L in β Th-patients have now been adopted by all transfusion centers worldwide². In addition to the cardiovascular and hematologic benefits gained in this manner, it is especially beneficial to bone health as well; hypoxia stops to exist and the erythropoietic marrow stops being stimulated and expanding^{4-6,8}.
- II. Regular transfusions unfortunately lead to excessive iron deposition in the tissues, the cytotoxic effect of which is responsible for the development of serious to fatal complications in multiple organs^{70,71}. As early as the 1970's, chelating agents have provided a solution to this problem, with the introduction of the subcutaneously administered chelating factor deferoxamine. From 1990's onwards, two other orally administered agents have been introduced, deferiprone and deferasirox, so that now, depending on the particularities of each patient, a completely individualized chelation regimen can be implemented, entirely eliminating -in the majority of patients- unnecessary iron from tissues⁷⁰⁻⁷². The removal of excess iron from the skeleton itself has an extremely favorable effect on bone health. In addition, successful chelation leads to improvement in the functioning of the endocrine glands and adequate production of hormones that have a positive effect on bone metabolism. Trials and case reports exist from recent years pertaining even to full reversal of endocrine disorders such as hypothyroidism, hypoparathyroidism, hypogonadism and diabetes mellitus^{73,74}. The above observations have also been confirmed by a recent study that came to light in 2014, conducted by Casale M. et al, who studied the positive effects of deferasirox on the endocrine

glands and bone. Among other things, they confirmed a significant improvement in the BMD of the spine during administration of the drug ($p < 0.001$), a trend of improvement in the BMD of the hip ($p = 0.2$), as well as a reduction in the percentage of patients with osteoporosis of the lumbar spine. In addition, they documented that these bone effects were independent of the effects of any anti-osteoporosis treatment and were due exclusively to successful long-term chelation⁶⁹. On the other hand, as already noted, the excessive or incorrect use of chelating agents may significantly worsen bone metabolism and BMD. On the basis of the above, proper and individualized use of chelating preparations following assessment of: a) the clinical history of each patient, b) serum ferritin levels and c) measurements of tissue-iron, using newer specialized MRI-techniques, is the strategy proposed today, initially to improve and then to maintain, both the skeleton as well as the whole body of β Th-patients in the best possible condition^{5,24,29-32}.

- III. Prevention and treatment of other co-morbidities of patients with β Th also constitute a key factor, which is favorable to their skeletal health. Complications such as heart failure and severe liver dysfunction represent difficult-to-control entities which, in addition to being able to become life-threatening, have an adverse effect on the skeleton. Monitoring of patients by a cardiologist and a hepatologist is deemed necessary^{4,5}. Furthermore, almost all the hormonal deficiencies gradually lead to major bone damage. Therefore, successful regulation of these endocrine disorders is extremely beneficial in TBD, among other things. Special mention should be made of the most common endocrine complication found in beta-thalassemia, hypogonadism. It has been established that hypogonadism and its treatment through appropriate hormone replacement therapy (HRT), have a significant impact on BMD and the incidence of fractures in the general population⁷⁵. Similar trials in β Th-patients have demonstrated the benefits of HRT in improving bone mass, even though there are still no clear data on the effect on the frequency of fractures³⁹. Yet another characteristic of TBD is that β Th-sufferers with hypogonadism, although greatly benefited by the administration of HRT, fail to improve, as regards BMD, to the same extent as non-thalassemic patients with hypogonadism receiving the same treatment^{39,76}. Furthermore, the gonadal hormones must be used with special caution in β Th; due to the hepatic, cardiac and vascular disorders, the safer transdermally administered forms of androgen and estrogen are preferable to the oral or intramuscular forms⁷⁷. Also, the dosage of these hormones should be the lowest possible for each patient, so as to ensure a positive impact on the bones while minimizing the risk of further complications. Finally, the indications for, and length of, therapy must also be carefully assessed and periodically adjusted, as these vary by age, the presence

of thrombophilia, splenectomy, diabetes and various other pathological states^{4,78}. As regards the inadequate functioning of the GH/IGF-1 axis, we should mention that the effects of administering the recombinant-GH in adult populations of β Th-patients have not been studied thus far at the TBD level and, consequently, the hormone must be administered only on the basis of the indications it has in childhood.

C. Management of hypercalciuria

As previously mentioned, hypercalciuria occurs in about one-third of β Th-patients. In order to control this disorder, the implementation of dietary measures that reduce the excretion of calcium in the urine (see above) and pertain to good hydration is initially recommended. Especially as regards the intake of calcium, this should not be entirely restricted, but must range between 800-1000 mg of calcium daily, preferably with food, and evenly distributed with meals. Without calcium supplementation bone health is bound to worsen significantly while, in addition, a mechanism is triggered for the development of hyperoxaluria and the formation of oxalate stones⁷⁹. At the same time, the intake of vitamin-D should also not be limited, but should be ingested at a sufficient dose to maintain 25OHD levels at the lower end of the normal range⁸⁰.

When the above measures are not sufficient and the hypercalciuria persists, it is recommended to commence treatment with thiazide diuretics (TZD). In clinical practice, hydrochlorothiazide and bendroflumethiazide are used today, administered as monotherapy or in combination with potassium-sparing diuretics. In general, as regards TZD, it is recommended that these be administered only when absolutely indicated, and in the lowest possible doses. Beta-thalassemia patients often maintain normal or borderline-low arterial blood pressure readings, and the risk of developing hypotension is significant. Also, it is necessary to evaluate sodium and potassium levels after the first 2-3 weeks of starting the medication, as the likelihood of hyponatremia and hypokalemia is not negligible^{81,82}. It should also be noted that when hypercalciuria co-exists with significant bone disease, the initial administration of TZD is recommended since, even if the hypercalciuria is regulated through dietary measures, the bone disease is not bound to improve⁸³. Finally, if bisphosphonates or denosumab be administered as an anti-osteoporosis therapy in a patient with hypercalciuria who also suffers from severe TBD, the hypercalciuria is also expected to improve to a certain extent⁸⁴.

D. Vitamins and minerals

Just as in the general population, so too with beta-thalassemia, adequate vitamin-D levels and adequate calcium intake significantly contribute to maintaining and improving bone health and preventing the development of osteomalacia. In β Th-patients, 25OHD levels are very often lower than normal, thus requiring laboratory testing

every 6-12 months and exogenous supplementation when indicated^{4,5}. Although sufficient data do not exist to determine the ideal 25OHD levels in β Th, recent trials have demonstrated positive non-linear correlation between the concentrations of 25OHD and BMD (Z-Score), with a plateau at concentrations of 25OHD above 15 ng/ml. These levels are therefore considered to be at minimum necessary in order for the BMD not to deteriorate⁸. In other trials, medical correction is recommended for levels below 20 ng/ml, with a proposed dose of vitamin-D3 of 2,000 IU p.o daily for a period of 8 weeks, and laboratory re-evaluation for possible continuation of administration. In order to maintain satisfactory levels, a daily intake of 800-1,000 IU of vitamin-D2 or D3 is recommended, especially in regions with decreased sunlight. Regimens with 50,000 IU of vitamin-D2 orally administered weekly or monthly have also been implemented for correction of deficiencies or maintenance of satisfactory levels respectively⁵⁴. Based on the above, in general, the recommended daily intake of vitamin-D does not seem to vary significantly from what is recommended for each age group in the healthy population^{4,5}. But, in β Th we must be more cautious, mainly due to the frequent hypercalciuria that is identified in patients. For the same reason, it is desirable for the level of 25OHD to be at approximately 30 ng/ml and not to exceed this by too much^{4,5,66}. It should be noted finally that, from all the relevant preparations, cholecalciferol should generally be preferred, while on the background of developing complications from other organs (hyperparathyroidism, liver dysfunction, kidney disorders) careful use of the more specific vitamin-D metabolites such as alfacalcidol and calcitriol may be required⁸⁵.

As regards calcium intake, the treating physician must decide, again guided by the pathophysiology and comorbidity of β Th. In general, patients should be encouraged to receive sufficient quantities of the mineral, according to the guidelines for the healthy population and mainly through meals, both because of the hypercalciuria as well as the potential coexistence of injury to the endothelium of the vessels due to iron deposits. If, however, the calcium must be administered through supplementation, low doses are preferable, or splitting of higher doses in 2-3 individual doses over a 24-hour period, in addition to the foregoing, for optimal absorption by the gastrointestinal tract. Although there is no official contraindication for the administration of calcium through supplements, even today some researchers point to a possible correlation of its administration with cardiovascular events⁸⁶. At the same time, many other trials have failed to demonstrate any such correlation^{87,88}.

Furthermore, in β Th-patients deficiencies in other minerals, such as magnesium, zinc and copper are also observed. There is causal speculation pertaining to the involvement of malabsorption from the gastrointestinal tract and/or of the action of chelating agents. Consequently, both periodic hematological testing and correction of low mineral levels are deemed absolutely essential⁸⁹. The role of magnesium

in bone metabolism is fundamental. Zinc is another mineral that significantly affects bone health. Indeed, correction of zinc deficits has been shown to contribute significantly to improvement in BMD, but also to the metabolism of glucose in β Th-patients^{89,90}. Therefore, although no specific instructions exist to date for determining the ideal daily dose and frequency of administration of the above supplements, physicians should be encouraged to monitor their patients periodically and to take action whenever low levels are identified. Finally, it is recommended that vitamin-C be taken in sufficient quantities, through food or even with additional supplements, if a lack is identified^{66,67}.

E. Anti-osteoporosis medication

- (i) **CALCITONIN.** Calcitonin is one of the most well studied anti-osteoporosis drugs that have been administered to β Th-patients. It was widely used during earlier years without, however, leading to truly remarkable results. Nevertheless, it seemed to restrict bone loss to a certain extent, while its contribution to the reduction of bone pain due to microfractures was also significant. In subsequent years, with the advent of bisphosphonates, the use of calcitonin was practically abandoned^{4,2,4,5,17}.
- (ii) **BISPHOSPHONATES.** Bisphosphonates (BSPs) are potent inhibitors of the bone resorption, drastically reducing the induction, proliferation, differentiation and the lifespan of osteoclasts¹⁶. Over the past two decades, many trials have been published on the subject of assessing the behavior and the effectiveness of the various types of BSPs on the TBD⁹¹⁻¹⁰⁷. Out of these, it has been concluded, judging by the final increase in BMD, that the most effective category is that of the intravenously administered amino-bisphosphonates, represented by zoledronic acid (ZA), pamidronate, and neridronate (Table 1). Administration of these specific preparations has led to increases in BMD of up to 50% in the lumbar spine, and up to 40% in other locations of the skeleton⁹⁷⁻¹⁰¹. Out of the substances administered orally, the most favorable therapeutic results are observed with alendronate^{92,102,106}. Unfortunately, in most of these trials, the number of participants was quite small, their heterogeneity really great (in terms of age, gender, chelation regimen and co-morbidity) and the time they were followed was brief. What has become clear thus far is that, in β Th-patients greater resistance develops to BSPs than that which occurs at least in postmenopausal osteoporosis. To a certain extent, the multifactorial etiopathogenesis of the TBD is definitely responsible for this. For this reason, it has been proposed by many experts that in β Th-individuals, BSPs be administered at higher doses than the conventional ones used. With the exception of alendronate, which seems to increase BMD to a sufficient extent at the usual oral dose of 70 mg/week, more aggressive protocols than the "classic" ones have been tested in terms of intravenously

administered amino-bisphosphonates. In particular, ZA -probably the most effective bisphosphonate for TBD- has shown greater efficacy at doses of 4 mg/3months and 1 mg/3months, and especially during the first year of administration, with a significant reduction of biochemical bone turnover markers (BTMs)⁹⁷⁻¹⁰¹, but also with improvement in musculoskeletal pain when this was assessed through the use of special questionnaires^{99,100}. Pamidronate has been administered in monthly doses of 15, 30, 60 mg and 1 mg/Kg for a total of 1 to 3 years, also with satisfactory results as regards increasing BMD in the spine and hip, and a reduction in BTMs^{95,103-105}. More recently, the intravenous administration of neridronate has also yielded satisfactory results in increasing BMD and reducing pain¹⁰⁷ while, finally, it is noted that both intravenous and intramuscular administration of clodronate has proven practically ineffective^{92,94}.

It should be noted that, as regards the use of BSPs in TBD, no significant adverse effects have been described in the relevant trials published, at the dosages and intervals that have been utilized to date. However, the potential risk of atypical hip fractures or osteonecrosis of the jaw with long-term use of these drugs -due to a reduction in bone remodeling- has not yet been sufficiently investigated^{4,5,91}. Moreover, though a causal relationship cannot be proven, two reported cases of the development of atypical hip fractures in β Th-patients who had previously received BSPs have already been published^{108,109}. Furthermore, in 2014, three cases were announced of β Th-patients who had previously taken p.o bisphosphonates, presenting with osteonecrosis of the jaw diagnosed at stage zero¹¹⁰. Therefore, the administration of BSPs must be evaluated individually for each patient and must be determined very carefully. Since there are no formal specific guidelines, neither as regards the suggested length of administration nor the dosage regimens, many researchers believe that the most logical approach is for BSPs to initially be administered on the basis of the thoroughly tested standards pertaining to postmenopausal osteoporosis: 5 years for oral and 3 years for i.v. BSPs (with the safest dose being ZA in 5 mg/year), and for the results be re-assessed at regular intervals⁹¹. Unfortunately, evaluation data on the effect of BSPs in preventing fractures in TBD, have not yet been published. The main reasons for this include the small number of participants and the short monitoring period in the trials carried out to date. Finally, since the anti-osteoporosis drugs are administered at significantly younger ages in TBD than in the general population, further individualized data will need to be taken into account as regards their use in each patient such as, for example, family planning, as BSPs remain in the body for long periods of time, and we are unfamiliar with the effects that they may have on the pregnancy and fetus^{111,112}.

- (iii) **DENOSUMAB.** One of the key-mechanisms of the pathogenesis of the TBD is the increased sRANKL/OPG ratio, resulting in excessive heightening of the osteoclasts function. It has even been noted that,

Table 1. Interventional studies on the administration of bisphosphonates for TBD since 2002^{92,94,95,98-107}. (M=Males, F=Females, IM=Intramuscular, m=month(s), d=days, w=week(s), ZA=Zoledronic Acid, vs=versus, Sign.=Significant, NS=non-significant, FN=Femoral neck, LS=Lumbar spine, BMD=Bone Mineral Density, Cal=Calcium, Vit.D=Vitamin-D).

AUTHOR YEAR	STUDY DESIGN	INTERVENTION	Duration (months)	Number & Sex of participants	Age (years)
Morabito et al, 2002 ⁹²	Randomized, parallel arm, placebo-controlled	Alendronate 10 mg/d (A) vs IM Clodronate 100 mg/10d (B) vs placebo (C)	24	25 6M/19F	26.6 ± 7.1
Results: (A) ↑ 2.8% LS BMD, ↑ 5.64% FN BMD All Sign. vs placebo, (B) ↔ LS and FN BMD vs placebo and baseline					
Pennisi et al, 2003 ⁹⁴	Randomized, controlled	Clodronate 300 mg/3w vs Cal+Vit.D	24	30 30M	27.44 ± 3.28
Results: ↔ LS and FN BMD T-Score NS vs baseline / p<0.01 vs Cal+Vit. D					
Voskaridou et al, 2003 ⁹⁵	Non-randomized, non-controlled, parallel arm	Pamidronate 30 mg/m (A) vs Pamidronate 60 mg/m (B)	12	26 6M/20F	35.5
Results: (A) and (B) ↑ LS BMD Z-Score / ↔ FN and Forearm BMD Z-Score vs baseline					
Gilfillan et al, 2006 ⁹⁸	Randomized, double-blind, placebo-controlled	ZA 4 mg/3m vs placebo	24	23 17M/6F	(ZA) 27.8 (18.2-40.4) (pl) 27.7 (18.9-43.7)
Results: ↑ 10.2% LS BMD / ↑ 5.2% FN BMD / ↑ 6% Tot.Hip BMD / ↑ 4.6% Tot.BMD - All Sign. vs placebo					
Voskaridou et al, 2006 ⁹⁹	Randomized, parallel arm, placebo-controlled	ZA 4 mg/6m (A) vs ZA 4 mg/3m (B) vs placebo (C)	12	66 22M/44F	(A) 44.1 ± 11.7 (B) 42.6 ± 10.7 (C) 44.9 ± 10.7
Results: (A) ↑ 5.8% LS BMD, ↑ 4.8% FN BMD NS vs placebo, (B) ↑ 15.2% LS BMD, ↑ 11.3% FN BMD Sign. vs placebo					
Otrock et al, 2006 ¹⁰⁰	Open label, non-randomized, controlled	ZA 4 mg/3m (A) vs observation (B)	12	28 18M/10F	(A) 22.72 ± 5.85 (B) 19.10 ± 4.07
Results: (A) ↑ LS and FN and Tot.Hip BMD Z-Score vs baseline, (B) ↔ LS and FN and Tot.Hip BMD Z-Score vs baseline					
Perifanis et al, 2007 ¹⁰¹	Single arm, non-controlled	ZA 1 mg/3m vs placebo	12	29 13M/16F	27.2 ± 7.3
Results: ↑ LS BMD T-Score vs baseline					
Skordis et al, 2008 ¹⁰²	Randomized, parallel arm, non-controlled	Alendronate 70 mg/w (A) vs Pamidronate 90 mg/m (B)	24	53 22M/31F	(A) 33.3 (20-47) (B) 34.4 (25-50)
Results: (A) ↔ LS and FN BMD Z-Score vs baseline, (B) ↑ LS and FN BMD Z-Score vs baseline					
Patiroglu et al, 2008 ¹⁰³	Single arm, non-controlled	Pamidronate 15 mg/3m	12	23 11M/12F	7-14
Results: ↑ FN BMD Sign vs baseline					
Leung et al, 2009 ¹⁰⁴	Non-randomized, parallel arm, controlled	Pamidronate (A) vs Cal+Alfacalcidol (B) vs Observation (C)	36	39 16M/23F	(A) 18.9 (17.5-22.4) (B) 22.2 (19.0-27.8) (C) 17.7 (16.3-21.3)
Results (Z-Scores): (A) ↑ LS & FN BMD vs baseline, (B) ↔ LS & FN BMD vs baseline, (C) ↔ LS & ↓ FN BMD vs baseline					
Chatterjee et al, 2012 ¹⁰⁵	Non-randomized, controlled	Pamidronate 1 mg/Kg/m (A) vs observation (B)	36	34 18M/16F	17-43
Results: (A) ↑ LS and Tot.Hip BMD Z-Score vs baseline, (B) ↔ LS and Tot.Hip BMD Z-Score vs baseline					
Shirani et al, 2012 ¹⁰⁶	Single arm, Non-controlled	Aledronate 10 mg/d	12	120 57M/69F	33 (20-50)
Results: ↑ LS and FN BMD T-Score vs baseline					
Forni et al, 2012 ¹⁰⁷	Randomized, open-label, controlled	Neridronate 100 mg/90d vs Cal+Vit.D	12	118 51M/67F	32.8 ± 8.1
Results: ↑ LS and FN and Tot. Hip BMD vs Cal+Vit. D					

despite the successful efforts in normalizing hemoglobin levels, appropriate HRT and substitution with thyroxine or GH, satisfactory control of diabetes and effective chelation, the sRANKL/OPG ratio continues to remain at pathologically high levels^{39,113,114}. Denosumab, a monoclonal anti-RANKL antibody that inhibits the binding of RANKL to RANK, would theoretically significantly correct this specific abnormality and would drastically reduce bone resorption¹¹⁵. The first non-controlled study on the effect of denosumab in TBD was published in 2014, with very encouraging results indeed¹¹⁶. In that trial, 30 patients with TBD aged 17 to 32, received the drug subcutaneously at the usual dose of 60 mg/6 months and, after 12 months, changes in BMD were evaluated, as were BTMs. A significant improvement in BMD of the femoral neck of 6.0%, an even greater improvement in the lumbar spine of 9.2%, and a considerable reduction of bone resorption markers were observed. Side effects were mild and mostly pertained to pain in the spine and the extremities (12%), as well as the presence of nausea (10%). The development of hypocalcemia in 7% of patients was mild and asymptomatic. Of course, the frequent presence of hypoparathyroidism and renal disorders in β Th requires attention to this specific matter.

The positive results of this trial have prompted the same researchers to launch, in 2017, a new randomized, open labeled, parallel assignment, interventional clinical phase-III trial to compare denosumab with ZA. A total of 10 patients were recruited in each group, and the initial results for changes in BMD and BTMs are expected to be announced in 2018. Finally, at the 22nd annual Congress of the European Hematology Association (EHA), which took place in Madrid in 2017, the initial results of the single-site, clinical trial by Voskaridou et al. were announced, which pertained to the administration of denosumab in 31 β Th-patients, at a dose of 60 mg/6 months for one year, compared with the administration of a placebo in 30 other β Th-patients. The results after the first two doses of the drug, here too revealed significant increases in the BMD in the denosumab group, both in the lumbar spine (6.02 \pm 5.30%) and the femoral neck, clearly superior to the placebo, and an excellent safety profile, with no significant side effects (*Voskaridou E. et al. Denosumab increases BMD in patients with thalassemia-major and osteoporosis: Results of a randomized, placebo-controlled, double-blind, phase-2B Clinical Trial - EHA22, June 2017 / Unpublished data*).

One observation that should be emphasized at this point is that, among the side effects of denosumab, a possible mild increase in the likelihood of the development of certain infections is mentioned, such as upper respiratory tract, urinary, skin and ear infections¹¹⁷. Since β Th-patients represent a group with a mildly increased susceptibility to infections^{118,119} and often have a positive history for splenectomy, heart disease, and various co-morbidities,

this observation should always be assessed by their treating physicians, without of course necessarily precluding the use of denosumab for TBD. Moreover, in the trials to date, there are no reports of a causal relationship between the drug with the development of infections in β Th-individuals.

(iv) TERIPARATIDE. Teriparatide is currently the only anabolic drug marketed for the treatment of osteoporosis. Nonetheless, its use in TBD has not yet been sufficiently studied, although it represents a tempting choice since very low BMD values are often observed in β Th-patients. Moreover, osteoblasts and bone formation are significantly affected in this hematologic disease⁹¹. On the other hand, due to its mechanism of action, the question remains as to whether or not this can significantly worsen any possible concomitant hypercalciuria. In addition, in quite a few transfused patients, foci of severe extramedullary hematopoiesis develop, which lead to serious pressure phenomena in the spine and require the implementation of radiotherapy. In these cases, the use of teriparatide is contraindicated¹⁶. To date, only three case reports have been published on the use of this drug in a corresponding number of β Th-patients with serious TBD and pathological fractures, and the truth is that, in all three, the results were very encouraging, with notable increases in BMD in the lumbar spine, the femoral neck and the total hip, at 12 and 18 months¹²⁰⁻¹²². Of course, more data are clearly required to substantiate both the efficacy and safety of teriparatide for its generalized use in severe TBD.

(v) STRONTIUM RANELATE. In terms of the effectiveness of strontium ranelate in TBD, only one randomized controlled trial was conducted, with relatively positive results, as an increase in BMD was observed -mainly in the spine- as well as normalization of BTMs, a reduction in sclerostin levels in the serum, and a reduction in musculoskeletal pain¹²³. Nevertheless, the drug practically ceased to represent a treatment option for TBD from the moment that new data arose pertaining to the safety of its administration, which were related to an increased risk of cardiovascular complications¹²⁴. These situations render it dangerous in practice for a patient-population with an increased prevalence of a variety of cardiovascular and thrombophilic problems. In any case, strontium ranelate is no longer available, as its manufacturers decided to cease supply of the drug treatment in August 2017.

(vi) NEW THERAPEUTIC AGENTS

Antibodies against sclerostin. Romosozumab is a monoclonal antibody that selectively binds to sclerostin and inhibits its action. It belongs to the anabolic drugs, as its administration significantly increases bone formation, and through this mechanism dramatically improves, in trials to date, the BMD in the spine and hip¹²⁵. Although its action in TBD has not yet been examined, scientific interest in its potential efficacy is

great, as significantly higher levels of sclerostin have been found in the serum of β Th-patients compared to the general population^{16,126,127}. Of course, the relatively recent reports of potentially increased incidence of cardiovascular episodes in clinical phase-III trials (“ARCH” and “BRIDGE”), draw attention for a potential future use of the drug in TBD.

Activin-A Inhibitors. Activins are members of the transforming growth factor- β (TGF- β) superfamily and carry out multiple biological actions in several tissues. Activin-A is a protein that is expressed particularly in the bones and it has been observed to be increased in the sera of patients with osteoporosis. In humans, it has been documented to promote osteoclastogenesis and differentiation of osteoclasts, while simultaneously inhibits differentiation and the action of osteoblasts^{128,129}. Based on these observations, agents have been developed which inhibit the effects of activin-A on the skeleton and are expected to improve bone status in individuals with osteoporosis. The implementation of this type of treatment in TBD has garnered great interest in recent years since, apart from the action of these drugs on the bone metabolism, a simultaneous significant increase in hemoglobin levels was identified in the preclinical trials on experimental animal models¹³⁰⁻¹³². The action on the hematopoietic tissue is carried out in the final stages of differentiation of red blood cells, correcting -to a certain extent- the ineffective erythropoiesis that occurs in β Th. In this manner, the need for transfusions, hyperplasia of the hemopoietic marrow, and hemosiderosis are reduced and quality of life is improved. On the level of clinical trials in β Th-patients, two separate molecules of activin-A inhibitors have already been administered, sotatercept and luspatercept, with very good results in terms of hematological data, but also with only mild side effects¹³³⁻¹³⁵. At this stage, research continues only with luspatercept, for which is underway, since 2016, an international multicenter randomized, double-blind phase-III trial (“BELIEVE” Clinical Trial) with a three years duration, with the participation of 300 β Th-patients. Of these, 200 receive the medicinal agent at varying doses subcutaneously every 3 weeks and the remaining 100 have been randomized in the placebo group. The first results are expected to be announced in 2018, and these will focus primarily on the hematologic data. However, some parameters which pertain to the effects of the drug on the bone metabolism, will also be tested¹³⁶.

Conclusion

β Th is a serious hematologic disorder with many accompanying complications, part of which is the early development of a particular type of bone disease with a multifactorial etiopathology. The treating physician is initially called upon to implement the necessary preventive measures in order to limit the unfavorable development

of skeletal problems and secondly, if required, to create a completely specialized and individualized treatment plan for each patient. Maintaining sufficient levels of hemoglobin, effective chelation, the most effective possible treatment of co-morbidities, proper nutrition, adequate calcium intake, correction of deficits of other minerals such as magnesium and zinc, strengthening of vitamin-D stores, management of hypercalciuria and appropriate exercise, now represent the cornerstones of a comprehensive preventive and therapeutic approach, even before the use of anti-osteoporosis drugs. Of course, with the most serious forms of TBD, the implementation of a more specialized medication regimen is strongly recommended. In such cases, the administration of BSPs was, until a few years ago, the only option for which numerous officially published data existed. Although BSPs still remain by far the most adequately studied medications in TBD, in recent years, more and more trials are coming to light on the use of new agents as well, with very promising results. Of these, very important information has been announced about denosumab, pertaining to satisfactory increases in BMD after 12 months of administration. The role of teriparatide should also be explored further. Finally, newer substances that are being developed, such as romosozumab and luspatercept, pique the interest and optimism of both doctors and patients. As the data to date are deemed to be encouraging and optimistic on the whole, the therapeutic approach of these patients is expected to be more multidimensional in the very near future.

References

1. Olivieri NF. The β -Thalassemias. *N Engl J Med* 1999;341(2):99-109.
2. Cappellini MD, Cohen A, Porter J, Taher A, Viprakasit V. Guidelines for the management of transfusion dependent thalassaemia (TDT), 3rd edition. Nicosia (CY): Thalassaemia International Federation; 2014.
3. Galanello R, Origa R. Beta-thalassemia. *Orphanet J Rare Dis* 2010; 5:11.
4. Toumba M, Skordis N. Osteoporosis syndrome in thalassaemia major: an Overview. *J Osteoporos* 2010;2010:537673.
5. Terpos E, Voskaridou E. Treatment options for thalassemia patients with osteoporosis. *Ann NY Acad Sci* 2010;1202:237-43.
6. Baldini M, Forti S, Marcon A, Olivieri FM, Orsatti A, Tampieri B et al. Endocrine and bone disease in appropriately treated adult patients with beta- thalassemia major. *Ann Hematol* 2010;89(12):1207-13.
7. Dresner Pollack R, Rachmilewitz E, Blumenfeld A, Idelson M, Goldfarb AW. Bone mineral metabolism in adults with beta-thalassemia major and intermedia. *Br J Hematol* 2000;111(3):902-7.
8. Vogiatzi MG, Macklin EA, Fung EB, Cheung AM, Vichinsky E, Olivieri N, et al. Bone disease in thalassemia: a frequent and still unresolved problem. *J Bone Miner Res* 2009;24(3):543-57.
9. Vogiatzi MG, Macklin EA, Fung EB, Vichinsky E, Olivieri N, Kwiatkowski J, et al. Prevalence of fractures among the Thalassemia syndromes in North America. *Bone* 2006;38(4):571-5.
10. Sutipornpalangkul W, Janechetsadatham Y, Sritanaratkul N, Harnroongroj T. Prevalence of fractures among Thais with thalassaemia syndromes. *Singapore Med J* 2010;51(10):817-21.

1. Fung EB, Harmatz PR, Milet M, Coates TD, Thompson AA, Ranalli M, et al. Fracture prevalence and relationship to endocrinopathy in iron overloaded patients with sickle cell disease and thalassemia. *Bone* 2008;43(1):162-8.
12. Origa R, Fiumana E, Gamberini MR, Armari S, Mottes M, Sangalli A, et al. Osteoporosis in β -Thalassemia: Clinical and genetic aspects. *Ann NY Acad Sci* 2005;1054(1):451-6.
13. Christoforidis A, Hatzipantelis E, Tsatra I, Kazantzidou E, Katzos G, Athanassiou-Metaxa M. Bone mineral density in children and young adults with β -thalassemia major conventionally treated. *Pediatr Blood Cancer* 2006;47(1):113-4.
14. Kyriakou A, Savva SC, Savvides I, Pangalou E, Ioannou YS, Christou S, et al. Gender differences in the prevalence and severity of bone disease in thalassaemia. *Pediatr Endocrinol Rev* 2008;6(Suppl. 1):116-22.
15. Ladis V, Raptou P, Rigatou E, Chouliaras G, Korres D, Kattamis C. Study of bone density by pQCT analysis in healthy adults and patients with B-Thalassemia major and intermedia. *Pediatr Endocrinol Rev* 2008;6(Suppl. 1):127-31.
16. Wong P, Fuller PJ, Gillespie MT, Milat F. Bone Disease in Thalassemia: A Molecular and Clinical Overview. *Endocr Rev* 2016;37(4):320-46.
17. Voskaridou E, Terpos E. New insights into the pathophysiology and management of osteoporosis in patients with β thalassaemia. *Br J Haematol* 2004;127(2):127-39.
18. Wonke B, Jensen C, Hanslip JJ, Prescott E, Lalloz M, Layton M, et al. Genetic and acquired predisposing factors and treatment of osteoporosis in thalassaemia major. *J Pediatr Endocrinol Metab* 1998;11(Suppl.3):795-801.
19. Skordis N, Efstathiou E, Kyriakou A, Toumba M. Hormonal dysregulation and bones in thalassaemia - an overview. *Pediatr Endocrinol Rev* 2008;6(Suppl. 1):107-15.
20. Vichinsky EP. The morbidity of bone disease in Thalassemia. *Ann NY Acad Sci* 1998;30(850):344-8.
21. Borgna-Pignatti C, Cappellini MD, De Stefano P, Del Vecchio GC, Forni GL, Gamberini MR, et al. Survival and complications in thalassemia. *Ann NY Acad Sci* 2005;1054:40-7.
22. Mahachoklertwattana P, Pootrakul P, Chuansumrit A, Choubtum L, Sriphrapradang A, Sirisriro R, et al. Association between bone mineral density and erythropoiesis in Thai children and adolescents with thalassemia syndromes. *J Bone Miner Metab* 2006;24(2):146-52.
23. Wong P, Fuller PJ, Gillespie MT, Kartsogiannis V, Strauss BJ, Bowden D, et al. Thalassemia bone disease: the association between nephrolithiasis, bone mineral density and fractures. *Osteoporos Int* 2013;24(7):1965-71.
24. Mahachoklertwattana P, Sirikulchayanonta V, Chuansumrit A, Karnsombat R, Choubtum L, Sriphrapradang A, et al. Bone histomorphometry in children and adolescents with β -thalassemia disease: iron-associated focal osteomalacia." *J Clin Endocrinol Metab* 2003;88(8):3966-72.
25. Zarjou A, Jeney V, Arosio P, Poli M, Zavaczki E, Balla G, et al. Ferritin ferroxidase activity: a potent inhibitor of osteogenesis. *J Bone Miner Res* 2010;25(1):164-72.
26. Yamasaki K, Hagiwara H. Excess iron inhibits osteoblast metabolism. *Toxicol Lett* 2009;191(2-3):211-5.
27. Yang Q, Jian J, Abramson SB, Huang X. Inhibitory effects of iron on bone morphogenetic protein 2-induced osteoblastogenesis. *J Bone Miner Res* 2011;26(6):1188-96.
28. Jia P, Xu YJ, Zhang ZL, Li K, Zhang W, Yang H. Ferric ion could facilitate osteoclast differentiation and bone resorption through the production of reactive oxygen species. *J Orthop Res* 2012;30(11):1843-52.
29. De Virgiliis S, Congia M, Frau F, Argioli F, Diana G, Cucca F, et al. Deferoxamine-induced growth retardation in patients with thalassemia major. *J Pediatr* 1988;113(4):661-9.
30. Naderi M, Sadeghi-Bojd S, Valeshabad AK, Jahantigh A, Alizadeh S, Dorgalaleh A, et al. A prospective study of tubular dysfunction in pediatric patients with beta thalassemia major receiving deferasirox. *Pediatr Hematol Oncol* 2013;30(8):748-54.
31. Quinn CT, Johnson VL, Kim HY, Trachtenberg F, Vogiatzi MG, Kwiatkowski JL, et al. Renal dysfunction in patients with thalassaemia. *Br J Haematol* 2011;153(1):111-7.
32. Wong P, Polkinghorne K, Kerr PG, Doery JC, Gillespie MT, Larmour I, et al. Deferasirox at therapeutic doses is associated with dose-dependent hypercalciuria. *Bone* 2016;85:55-8.
33. Chan YL, Pang LM, Chik KW, Cheng JC, Li CK. Patterns of bone diseases in transfusion-dependent homozygous thalassaemia major: predominance of osteoporosis and desferrioxamine-induced bone dysplasia. *Pediatr Radiol* 2002;32(7):492-7.
34. Olivieri NF, Koren G, Harris J, Khattak S, Freedman MH, Templeton DM, et al. Growth failure and bony changes induced by deferoxamine. *Am J Pediatr Hematol Oncol* 1992;14(1):48-56.
35. Morabito N, Gaudio A, Lasco A, Atteritano M, Pizzuolo MA, Cincotta M, et al. Osteoprotegerin and RANKL in the pathogenesis of thalassemia-induced osteoporosis: new pieces of the puzzle. *J Bone Miner Res* 2004;19(5):722-7.
36. Voskaridou E, Christoulas D, Xirakia C, Varvagiannis K, Boutsikas G, Bilalis A, et al. Serum Dickkopf-1 is increased and correlates with reduced bone mineral density in patients with thalassaemia-induced osteoporosis. Reduction post-zoledronic acid administration. *Haematologica* 2009;94(5):725-8.
37. Morabito N, Russo GT, Gaudio A, Lasco A, Catalano A, Morini E, et al. The "lively" cytokines network in beta-Thalassemia Major-related osteoporosis. *Bone* 2007;40(6):1588-94.
38. Borgna-Pignatti C, Rugolotto S, De Stefano P, Zhao H, Cappellini MD, Del Vecchio GC, et al. Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica* 2004;89(10):1187-93.
39. Anapliotou ML, Kastanias IT, Psara P, Evangelou EA, Liparaki M, Dimitriou P. The contribution of hypogonadism to the development of osteoporosis in thalassaemia major: new therapeutic approaches. *Clin Endocrinol (Oxf)* 1995;42(3):279-287.
40. Gamberini MR, De Sanctis V, Gilli G. Hypogonadism, diabetes mellitus, hypothyroidism, hypoparathyroidism: incidence and prevalence related to iron overload and chelation therapy in patients with thalassaemia major followed from 1980 to 2007 in the Ferrara centre. *Pediatr Endocrinol Rev* 2008;6(Suppl. 1):158-69.
41. Shamsirsaz AA, Bekheirnia MR, Kamgar M, Pourzahedgilani N, Bouzari N, Habibzadeh M, et al. Metabolic and endocrinologic complications in beta-thalassemia major: a multicenter study in Tehran. *BMC Endocr Disord* 2003;3(1):4.
42. Toumba M, Sergis A, Kanaris C, Skordis N. Endocrine complications in patients with thalassaemia major. *Pediatr Endocrinol Rev* 2007;5(2):642-8.
43. Soliman AT, El Banna N, Ansari BM. GH response to provocation and circulating IGF-I and IGF-binding protein-3 concentrations, the IGF-I generation test and clinical response to GH therapy in children with beta-thalassaemia. *Eur J Endocrinol* 1998;138(4):394-400.
44. Karamifar H, Karimi M, Amirhakimi G, Sharbatialaei M, De Sanctis V. Reduced insulin growth factor I concentrations in iron-overloaded beta thalassaemic patients with normal growth hormone secretion and liver function." *Pediatr Endocrinol Rev* 2004;2(Suppl.2):256-8.
45. De Sanctis V. Growth and puberty and its management in thalassaemia. *Horm Res* 2002;58(Suppl 1):72-9.
46. Poggi M, Pascucci C, Monti S, Pugliese P, Lauri C, Amodio G, et al. Prevalence of growth hormone deficiency in adult polytransfused-

- thalassemia patients and correlation with transfusional and chelation parameters. *J Endocrinol Invest* 2010;33(8):534-8.
47. Italian Working Group on Endocrine Complications in Non-Endocrine Diseases. Multicentre study on prevalence of endocrine complications in thalassaemia major. *Clin Endocrinol* 1995;42(6):581-6.
 48. Monge L, Pinach S, Caramellino L, Bertero MT, Dall'omo A, Carta Q. The possible role of autoimmunity in the pathogenesis diabetes in β -thalassaemia major. *Diabetes Metab* 2001;27(2 Pt 1):149-54.
 49. Lala R, Chiabotto P, Di Stefano M, Isaia GC, Garofalo F, Piga A. Bone density and metabolism in thalassaemia. *J Pediatr Endocrinol Metab* 1998;11(Suppl.3):785-90.
 50. Hamidieh AA, Moradbeag B, Pasha F, Jalili M, Hadjibabaie M, Keshavarznia M. High Prevalence of Hypoparathyroidism in patients with β -Thalassaemia Major. *Int J Hematol Oncol Stem Cell Res* 2009;3(3):17-20.
 51. Angelopoulos NG, Goula A, Rombopoulos G, Kaltzidou V, Katounda E, et al. Hypoparathyroidism in transfusion-dependent patients with β -thalassaemia. *J Bone Miner Metab* 2006;24(2):138-45.
 52. Vogiatzi MG, Macklin EA, Trachtenberg FL, Fung EB, Cheung AM, Vichinsky E, et al. Differences in the prevalence of growth, endocrine and vitamin D abnormalities among the various thalassaemia syndromes in North America. *Br J Haematol* 2009;146(5):546-56.
 53. Napoli N, Carmina E, Bucchieri S, Sferrazza C, Rini GB, Di Fede G. Low serum levels of 25-hydroxy vitamin D in adults affected by thalassaemia major or intermedia. *Bone* 2006;38(6):888-92.
 54. Soliman A, De Sanctis V, Yassin M. Vitamin D Status in Thalassaemia Major: an Update. *Mediterr J Hematol Infect Dis* 2013;5(1):e2013057.
 55. Mohkam M, Shamsian BS, Gharib A, Nariman S, Arzani MT. Early markers of renal dysfunction in patients with β -thalassaemia major. *Pediatr Nephrol* 2008;23(6):971-6.
 56. Koliakos G, Papachristou F, Koussi A, Perifanis V, Tsatra I, Souliou E, et al. Urine biochemical markers of early renal dysfunction are associated with iron overload in β -thalassaemia. *Clin Lab Hematol* 2003;25(2):105-9.
 57. Smolkin V, Halevy R, Levin C, Mines M, Sakran W, Ilia K, et al. Renal function in children with β -thalassaemia major and thalassaemia intermedia. *Pediatr Nephrol* 2008;23(10):1847-51.
 58. Sadeghi-Bojd S, Hashemi M, Karimi M. Renal tubular function in patients with β -thalassaemia major in Zahedan, southeast Iran. *Singapore Med J* 2008;49(5):410-2.
 59. Schouten BJ, Hunt PJ, Livesey JH, Frampton CM, Soule SG. FGF23 elevation and hypophosphatemia after intravenous iron polymaltose: a prospective study. *J Clin Endocrinol Metab* 2009;94(7):2332-7.
 60. Milat F, Wong P, Fuller PJ, Johnstone L, Kerr PG, Doery JC, et al. A case of hypophosphatemic osteomalacia secondary to deferasirox therapy. *J Bone Miner Res* 2012;27(1):219-22.
 61. Moulas A, Challa A, Chaliasos N, Lapatsanis PD. Vitamin D metabolites (25-hydroxyvitamin D, 24,25-dihydroxyvitamin D and 1,25-dihydroxyvitamin D) and osteocalcin in β -thalassaemia. *Acta Paediatr* 1997;86(6):594-9.
 62. Fung EB, Xu Y, Trachtenberg F, Odame I, Kwiatkowski JL, Neufeld EJ, et al. Inadequate dietary intake in patients with thalassaemia. *J Acad Nutr Diet* 2012;112(7):980-90.
 63. Valizadeh N, Farrokhi F, Alinejad V, Said Mardani S, Valizadeh N, Hejazi S, et al. Bone density in transfusion dependent thalassaemia patients in Urmia, Iran. *Iran J Ped Hematol Oncol* 2014;4(2):68-71.
 64. Shivani S, Douglas PK. Smoking, Alcohol and Bone. In: Hollich MF, Nieves JW. *Nutrition and Bone Health*, Edition: 2nd. New York: Springer; 2015. P. 489-504.
 65. Rosen CJ, Compston JE, Lian JB. *Primer on the metabolic bone diseases and disorders of mineral metabolism*, Edition: 7th. Washington, DC: ASBMR; 2008.
 66. Angelucci E, Barosi G, Camaschella C, Cappellini MD, Cazzola M, Galanello R, et al. Italian Society of Hematology guidelines for the management of iron overload in thalassaemia major and related disorders. *Haematologica* 2008;93(5):741-52.
 67. Yardumian A, Telfer P, Shah F, Ryan K, Darlison MW, Miller E, Constantinou G. *Standards for the Clinical Care of Children and Adults with Thalassaemia in the UK*, Edition: 3rd. London: United Kingdom Thalassaemia Society; 2016.
 68. Perrotta S, Cappellini MD, Bertoldo F, Servedio V, Iolascon G, D'Agruma L, et al. Osteoporosis in β -thalassaemia major patients: analysis of the genetic background. *Br J Haematol* 2000;111(2):461-6.
 69. Casale M, Citarella S, Filosa A, De Michele E, Palmieri F, Ragozzino A, et al. Endocrine function and bone disease during long-term chelation therapy with deferasirox in patients with β -thalassaemia major. *Am J Hematol* 2014;89(12):1102-6.
 70. Kwiatkowski JL. Management of transfusional iron overload - differential properties and efficacy of iron chelating agents. *J Blood Med* 2011;2:135-49.
 71. Cappellini MD, Pattoneri P. Oral iron chelators. *Ann Rev Med* 2009;60:25-38.
 72. Mobarra N, Shanaki M, Ehteram H, Nasiri H, Sahmani M, Saeidi M, et al. A Review on Iron Chelators in Treatment of Iron Overload Syndromes. *Int J Hematol Oncol Stem Cell Res* 2016;10(4):239-47.
 73. Farmaki K, Angelopoulos N, Anagnostopoulos G, Goula A, Pappa C. Thyroid Function in Thalassaemic Patients Treated with Combined Chelation Therapy. *Blood* 2006;108:1592.
 74. Farmaki K, Tzoumari I, Pappa C, Chouliaras G, Berdoukas V. Normalisation of total body iron load with very intensive combined chelation reverses cardiac and endocrine complications of thalassaemia major. *Br J Haematol* 2010;148(3):466-75.
 75. Cauley JA, Seeley DG, Ensrud K, Ettinger B, Black D, Cummings SR. Estrogen replacement therapy and fractures in older women. Study of Osteoporotic Fractures Research Group. *Ann Intern Med* 1995;122(1):9-16.
 76. Chatterjee R, Katz M, Bajoria R. Use of hormone replacement therapy for correction of high turnover bone disease in hypogonadal β -Thalassaemia major patients presenting with osteoporosis: comparison with idiopathic premature ovarian failure. *Hemoglobin* 2011;35(5-6):653-68.
 77. L'hermite M, Simoncini T, Fuller S, Genazzani AR. Could transdermal estradiol + progesterone be a safer postmenopausal HRT? A review. *Maturitas* 2008;60(3-4):185-201.
 78. Cappellini MD, Poggiali E, Taher AT, Musallam KM. Hypercoagulability in β -thalassaemia: a status quo. *Expert Rev Hematol* 2012;5(5):505-11.
 79. Curhan GC, Willett WC, Speizer FE, Spiegelman D, Stampfer MJ. Comparison of dietary calcium with supplemental calcium and other nutrients as factors affecting the risk for kidney stones in women. *Ann Intern Med* 1997;126(7):497-504.
 80. Han H, Segal AM, Seifter JL, Dwyer JT. Nutritional Management of Kidney Stones (Nephrolithiasis). *Clin Nutr Res* 2015;4(3):137-52.
 81. Moreira Guimaraes Penido MG, de Sousa Tavares M, Campos Linhares M, Silva Barbosa AC, Cunha M. Longitudinal study of bone mineral density in children with idiopathic hypercalciuria. *Pediatr Nephrol* 2012;27(1):123-30.
 82. Kruse C, Eiken P, Vestergaard P. Continuous and long-term treatment is more important than dosage for the protective effect of thiazide use on bone metabolism and fracture risk. *J Intern Med*

- 2016;279(1):110-22.
83. Schwadere AI, Srivastava T, Schueller L, Cronin R, Mahan JD, Hains D. Dietary modifications alone do not improve bone mineral density in children with idiopathic hypercalciuria. *Clin Nephrol* 2011;76(5):341-7.
 84. Arrabal-Polo MA, Arias-Santiago S, de Haro-Munoz T, Lopez-Ruiz A, Orgaz-Molina J, Gonzalez-Torres S, et al. Effects of aminobisphosphonates and thiazides in patients with osteopenia/osteoporosis, hypercalciuria, and recurring renal calcium lithiasis. *Urology* 2013;81(4):731-7.
 85. Brandi ML. Indications on the use of vitamin D and vitamin D metabolites in clinical phenotypes *Clin Cases Miner Bone Metab* 2010;7(3):243-50.
 86. Bolland MJ, Grey A, Avenell A, Gamble GD, Reid IR. Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women's Health Initiative limited access dataset and meta-analysis *BMJ* 2011;19;342:d2040.
 87. Lewis JR, Calver J, Zhu K, Flicker L, Prince RL. Calcium supplementation and the risks of atherosclerotic vascular disease in older women: results of a 5-year RCT and a 4.5-year follow-up. *J Bone Miner Res* 2011;26(1):35-41.
 88. Paik JM, Curhan GC, Sun Q, Rexrode KM, Manson JE, Rimm EB, et al. Calcium supplement intake and risk of cardiovascular disease in women. *Osteoporos Int* 2014;25(8):2047-56.
 89. Ozturk Z, Genc GE, Gumuslu S. Minerals in thalassaemia major patients: An overview. *J Trace Elem Med Biol* 2017;41:1-9.
 90. Fung EB, Kwiatkowski JL, Huang JN, Gildengorin G, King JC, Vichinsky EP. Zinc supplementation improves bone density in patients with thalassaemia: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr* 2013;98(4):960-71.
 91. Dede AD, Trovas G, Chronopoulos E, Triantafyllopoulos IK, Dontas I, Papaioannou N, et al. Thalassaemia-associated osteoporosis: a systematic review on treatment and brief overview of the disease. *Osteoporos Int* 2016;27(12):3409-25.
 92. Morabito N, Lasco A, Gaudio A, Crisafulli A, Di Pietro C, Meo A, et al. Bisphosphonates in the treatment of thalassaemia-induced osteoporosis. *Osteoporos Int* 2002;13(8):644-9.
 93. Viereck V, Emons G, Lauck V, Frosch KH, Blaschke S, Grundker C, et al. Bisphosphonates pamidronate and zoledronic acid stimulate osteoprotegerin production by primary human osteoblasts. *Biochem Biophys Res Commun* 2002;291(3):680-6.
 94. Pennisi P, Pizzarelli G, Spina M, Riccobene S, Fiore CE. Quantitative ultrasound of bone and clodronate effects in thalassaemia-induced osteoporosis. *J Bone Miner Metab* 2003;21(6):402-8.
 95. Voskaridou E, Terpos E, Spina G, Palermos J, Rahemtulla A, Loutradi A, et al. Pamidronate is an effective treatment for osteoporosis in patients with beta-thalassaemia. *Br J Haematol* 2003;123(4):730-7.
 96. Pan B, Farrugia AN, To LB, Findlay DM, Green J, Lynch K, et al. The nitrogen-containing bisphosphonate, zoledronic acid, influences RANKL expression in human osteoblast-like cells by activating TNF-alpha converting enzyme (TACE). *J Bone Miner Res* 2004;19(1):147-54.
 97. Perifanis V, Vyzantiadis T, Vakalopoulou S, Tziomalos K, Garipidou V, Athanassiou-Metaxa M, et al. Treatment of beta-thalassaemia-associated osteoporosis with zoledronic acid. *Br J Haematol* 2004;125(1):91-2.
 98. Gilfillan CP, Strauss BJ, Rodda CP, Bowden DK, Kean AM, Obaid M, et al. A randomized, double-blind, placebo-controlled trial of intravenous zoledronic acid in the treatment of thalassaemia-associated osteopenia. *Calcif Tissue Int* 2006;79(3):138-44.
 99. Voskaridou E, Anagnostopoulos A, Konstantopoulos K, Stoupa E, Spyropoulou E, Kiamouris C, et al. Zoledronic acid for the treatment of osteoporosis in patients with beta-thalassaemia: results from a single-center, randomized, placebo-controlled trial." *Haematologica* 2006;91(9):1193-202.
 100. O'trock ZK, Azar ST, Shamseddeen WA, Habr D, Inati A, Koussa S, et al. Intravenous zoledronic acid treatment in thalassaemia-induced osteoporosis: results of a phase II clinical trial. *Ann Hematol* 2006;85(9):605-9.
 101. Perifanis V, Vyzantiadis T, Tziomalos K, Vakalopoulou S, Garipidou V, Athanassiou-Metaxa M, et al. Effect of zoledronic acid on markers of bone turnover and mineral density in osteoporotic patients with beta-thalassaemia. *Ann Hematol* 2007;86(1):23-30.
 102. Skordis N, Ioannou YS, Kyriakou A, Savva SC, Efstathiou E, Savvides I, et al. Effect of bisphosphonate treatment on bone mineral density in patients with thalassaemia major. *Pediatr Endocrinol Rev* 2008;6(Suppl 1):144-8.
 103. Patiroğlu T, Torun YA, Kula M, Karakükçü M. Treatment of thalassaemia-induced osteoporosis with intermittent pamidronate infusions: two-year follow up. *Turk J Haematol* 2008;25(2):79-82.
 104. Leung TF, Chu Y, Lee V, Cheng FW, Leung WK, Shing MM, et al. Long-term effects of pamidronate in thalassaemic patients with severe bone mineral density deficits. *Hemoglobin* 2009;33(5):361-9.
 105. Chatterjee R, Shah FT, Davis BA, Byers M, Sooranna D, Bajoria R, et al. Prospective study of histomorphometry, biochemical bone markers and bone densitometric response to pamidronate in β -thalassaemia presenting with osteopenia-osteoporosis syndrome. *Br J Haematol* 2012;159(4):462-71.
 106. Shirani F, Iashkari M, Ahmadi AS, Arabi M, Asefi N. Evaluation of alendronate efficacy on bone mineral density in thalassaemic patients. *Int J Hematol Oncol Stem Cell Res* 2012;6(4):20-4.
 107. Forni GL, Perrotta S, Giusti A, Quarta G, Pitrolo L, Cappellini MD, et al. Neridronate improves bone mineral density and reduces back pain in β -thalassaemia patients with osteoporosis: results from a phase 2, randomized, parallel-arm, open-label study. *Br J Haematol* 2012;158(2):274-82.
 108. Giusti A, Barone A, Balocco M, Pinto V, Pilotto A, Forni GL. Atypical fractures of the femur and bisphosphonate therapy in patients with thalassaemia-associated osteoporosis: report of three cases. Poster presented at: 10th Cooley's Anemia Symposium; October 18-22, 2015; Chicago, IL.
 109. Lampropoulou-Adamidou K, Tournis S, Triantafyllopoulos IK. Atypical femoral fracture in a beta-thalassaemia major patient with previous bisphosphonate use: case report and a review of the literature. *J Musculoskelet Neuronal Interact* 2016;16(1):75-8.
 110. Chatterjee R, Bajoria R, Shah FT, Porter JB, Fedele S. High index of suspicion for early diagnosis of alendronate-induced stage zero osteonecrosis of jaw in thalassaemia major. *Br J Haematol* 2014;166(2):292-4.
 111. Stathopoulos IP, Liakou CG, Katsalira A, Trovas G, Lyritis GG, Papaioannou NA, et al. The use of bisphosphonates in women prior to or during pregnancy and lactation. *Hormones (Athens)* 2011;10(4):280-91.
 112. Levy S, Fayed I, Taguchi N, Han JY, Aiello J, Matsui D, et al. Pregnancy outcome following in utero exposure to bisphosphonates. *Bone* 2009;44(3):428-30.
 113. Carmina E, Di Fede G, Napoli N, Renda G, Vitale G, Lo Pinto C, et al. Hypogonadism and hormone replacement therapy on bone mass of adult women with thalassaemia major. *Calcif Tissue Int* 2004;74(1):68-71.
 114. Lasco A, Morabito N, Gaudio A, Buemi M, Wasniewska M, Frisina N. Effects of hormonal replacement therapy on bone metabolism in young adults with beta-thalassaemia major. *Osteoporos Int* 2001;

- 12(7):570-5.
115. Khosla S. Minireview: the OPG/RANKL/RANK system. *Endocrinology* 2001;142(12):5050-5.
 116. Yassin MA, Soliman AT, De Sanctis V, Abdelrahman MO, Aziz Bedair EM, AbdelGawad M. Effects of the anti-receptor activator of nuclear factor kappa B ligand denosumab on beta thalassemia major-induced osteoporosis. *Indian J Endocrinol Metab* 2014;18(4):546-51.
 117. CHMP Assessment report for Denosumab (Prolia). Doc.Ref.: EMA/21672/2010. Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Public_assessment_report/human/001120/WC500093529.pdf
 118. Ghaffari J, Abediankenari S, Nasehi M. Thalassemia and Immune System Dysfunction - Review Article. *Int J Curr Res* 2011;3(12):105-8.
 119. Farmakis D, Giakoumis A, Polymeropoulos E, Aessopos A. Pathogenetic aspects of immune deficiency associated with beta-thalassemia. *Med Sci Monit* 2003;9(1):RA19-22.
 120. Tournis S, Dede AD, Savvidis C, Triantafyllopoulos IK, Kattamis A, Papaioannou N. Effects of teriparatide retreatment in a patient with β -thalassemia major. *Transfusion* 2015;55(12):2905-10.
 121. Clima LN, Albu A, Lambrescu I, Barbu CG, Fica S. The response to teriparatide of a patient with β -thalassemia and multiple endocrine complications. *Endocrine Abstracts* 2015;37:EP243.
 122. Trotta A, Corrado A, Cantatore FP. Anabolic therapy of induced osteoporosis in beta-thalassaemia major: case report and literature review. *Reumatismo* 2010;62(2):119-26.
 123. Morabito N, Catalano A, Gaudio A, Morini E, Bruno LM, Basile G, et al. Effects of strontium ranelate on bone mass and bone turnover in women with thalassemia major-related osteoporosis. *J Bone Miner Metab* 2016;34(5):540-6.
 124. Reginster JY. Cardiac concerns associated with strontium ranelate. *Expert Opin Drug Saf* 2014;13:1209-13.
 125. McClung MR, Grauer A, Boonen S, Bolognese MA, Brown JP, Diez-Perez A, et al. Romosozumab in postmenopausal women with low bone mineral density. *N Engl J Med* 2014;370:412-20.
 126. Voskaridou E, Christoulas D, Plata E, Bratengeier C, Anastasilakis AD, Komninaka V, et al. High circulating sclerostin is present in patients with thalassemia-associated osteoporosis and correlates with bone mineral density. *Horm Metab Res* 2012;44(12):909-13.
 127. Saag KG, Petersen J, Brandi ML, Karaplis AC, Lorentzon M, Thomas T, et al. Romosozumab or Alendronate for Fracture Prevention in Women with Osteoporosis. *N Engl J Med* 2017;377(15):1417-27.
 128. Lotinun S, Pearsall RS, Home WC, Baron R. Activin receptor signaling: a potential therapeutic target for osteoporosis. *Curr Mol Pharmacol* 2012;5(2):195-204.
 129. Chantry AD, Heath D, Mulivor AW, Pearsall S, Baud'huin M, Coulton L, et al. Inhibiting activin-A signaling stimulates bone formation and prevents cancer-induced bone destruction in vivo. *J Bone Miner Res* 2010;25(12):2633-46.
 130. Langdon JM, Barkataki S, Berger AE, Cheadle C, Xue QL, Sung V, et al. RAP-011, an activin receptor ligand trap, increases hemoglobin concentration in hepcidin transgenic mice. *Am J Hematol* 2015;90(1):8-14.
 131. Fields SZ, Parshad S, Anne M, Raftopoulos H, Alexander MJ, Sherman ML, et al. Activin receptor antagonists for cancer-related anemia and bone disease. *Expert Opin Investig Drugs* 2013;22(1):87-101.
 132. Dussiot M, Maciel TT, Fricot A, Chartier C, Negre O, Velga J, et al. An activin receptor IIA ligand trap corrects ineffective erythropoiesis in β -thalassemia. *Nat Med* 2014;20(4):398-407.
 133. Cappellini MD, Porter J, Origa R, Forni GL, Laadem A, Galacteros F, et al. A Phase 2a, Open-Label, Dose-Finding Study To Determine The Safety and Tolerability Of Sotatercept (ACE-011) In Adults With Beta (β)-Thalassemia: Interim Results. *Blood* 2013;122:3448.
 134. Piga AG, Perrotta S, Melpignano A, Borgna-Pignatti, Gamberini MR, Voskaridou E, et al. Luspatercept decreases transfusion burden and liver iron concentration in regularly transfused adults with beta-thalassemia. Oral Presentation at: 21st Congress of European Hematology Association (EHA); June 9-12, 2016; Copenhagen, Denmark.
 135. Piga AG, Tartaglione I, Gamberini R, Voskaridou E, Melpignano A, Ricchie P, et al. Luspatercept Increases Hemoglobin, Decreases Transfusion Burden and Improves Iron Overload in Adults with Beta-Thalassemia. *Blood* 2016;128:851.
 136. Piga AG, Tartaglione I, Gamberini R, Voskaridou E, Melpignano A, Ricchie P, et al. Luspatercept increases hemoglobin and decreases transfusion burden in adults with β -thalassemia. Oral Presentation at: 22nd Congress of European Hematology Association (EHA); June 22-25, 2017; Madrid, Spain.