

Original Article

Association of Composite and Domain-Specific Intrinsic Capacity and Frailty Among Older Adults in India: Evidence from the Longitudinal Ageing Study in India (LASI)

Sudeep Mathew George^{#1}, Kritartha Kashyap^{#2}, Minakshi Dhar², Vasu Digra¹, Parul Bhutani², Pankhuri Saxena², Raj Kumar Tata¹, Bhabhor Kaminiben Narsinhbhai¹, Shreya Biswal¹, Pramod Kumar Mehta¹, Prasun Chatterjee³, Avinash Chakrawarty¹, Naveet Wig¹, Abhijith Rajaram Rao¹

¹Department of Geriatric Medicine, National Centre for Ageing, All India Institute of Medical Sciences, New Delhi, India

²Department of Geriatric Medicine, All India Institute of Medical Sciences, Rishikesh, India

³Department of Geriatric Medicine, Indraprastha Apollo Hospital, New Delhi, India

[#]Equal Contribution

Abstract

Objectives: Frailty and intrinsic capacity (IC), central constructs in the WHO healthy ageing framework, reflect vulnerability and resilience, respectively. Although both are linked to adverse outcomes, population-level evidence from India examining their relationship remains limited. This study explored the association between IC and frailty among older Indian adults. **Methods:** Data from 26,943 community-dwelling adults aged ≥ 60 years from the Longitudinal Ageing Study in India (LASI Wave 1) were analysed. Frailty was assessed using Fried's phenotype. IC was measured across cognition, locomotion, sensory, vitality, and mood domains, generating a composite score (0–10). Associations between IC and frailty status were examined using chi-square tests and multinomial logistic regression adjusted for sociodemographic factors and comorbidities. **Results:** Median age was 67 (63–72) years, and 48.4% were male. Frail individuals had lower median IC scores [6 (5–7)] than pre-frail and robust participants [7 (6–8); $p < 0.001$]. Higher IC scores were inversely associated with pre-frailty (RRR: 0.79, 95% CI: 0.75–0.84) and frailty (RRR: 0.67, 95% CI: 0.64–0.71). Poor cognition, locomotion, vitality, and mood were significantly associated with frailty, whereas sensory impairment was not. **Conclusion:** Lower IC is strongly associated with frailty among older Indian adults, supporting IC-based geriatric assessment for early identification of frailty progression.

Keywords: Intrinsic capacity, Frailty, Older adults, LASI, India

Introduction

Population ageing is a global phenomenon, with low- and middle-income countries like India experiencing especially rapid demographic shifts¹. As life expectancy increases, individuals often undergo multi-system dysregulation and a gradual decline in physiological reserves, which in turn increases their vulnerability to adverse outcomes, including disability, dependence, hospitalisation, and death². Consequently, maintaining functional ability and independence in older adults has become an important public health concern.

Frailty — a multidimensional geriatric syndrome

that reflects diminished resilience to stressors — has become a strong predictor of these adverse outcomes³. It has been operationalised in various ways. One widely

The authors have no conflict of interest.

Corresponding author: Abhijith Rajaram Rao, Department of Geriatric Medicine, National Centre for Ageing, All India Institute of Medical Sciences, New Delhi, 110029, India

E-mail: abhijith.rrao@gmail.com

Edited by: Jagadish Chhetri

Accepted 4 June 2026

used and validated model is Fried's frailty phenotype, which categorises individuals based on five clinical signs: unintentional weight loss (shrinking), weakness (grip strength), exhaustion, slowness (gait speed), and low physical activity. The presence of three or more criteria indicates frailty; one or two criteria suggest pre-frailty, and none indicates robustness⁴. This serves as a valuable clinical and public health framework for identifying older adults at high risk.

To complement this vulnerability-focused framework, the World Health Organization (WHO) has introduced the concept of intrinsic capacity (IC). IC is defined as the composite of all physical and mental abilities an individual can utilise at any given time and is typically conceptualized across five interconnected domains: cognition, locomotion, sensory ability, vitality (nutrition/energy), and psychological health^{5,6}. Unlike frailty, which emphasizes vulnerability, IC reflects a resilience-oriented approach, emphasising functional capacities that can be maintained or improved throughout the entire lifespan. IC is central to WHO's healthy ageing framework, as it concentrates on positive qualities and pathways across life⁷.

Although frailty and IC appear conceptually different, emerging evidence suggests they are closely related and may represent complementary perspectives of healthy ageing rather than competing constructs. Recent conceptual frameworks propose that declines in intrinsic capacity and the development of frailty occur along a continuum from resilience to vulnerability, with neither construct fully substituting for the other⁸⁻¹¹. Declines in IC have been shown to precede the onset of frailty and predict poor outcomes, including falls, disability, and mortality¹²⁻¹⁴. Longitudinal analyses demonstrate that worsening IC trajectories are strongly linked to the progression of frailty and disability, while higher IC acts as a safeguard against functional decline^{8,15}. Furthermore, impaired domains such as locomotion and vitality often serve as early indicators of frailty transitions^{16,17}. These findings support the idea that interventions aimed at preserving or restoring IC could delay or prevent frailty and its impacts^{8,13,18}.

In India, the Longitudinal Ageing Study in India (LASI, wave 1, 2017-18) provides a nationally representative dataset to examine both these concepts¹⁹. Recent LASI-based research has shown that only about one-quarter ($\approx 24.6\%$) of older adults (aged ≥ 60 years) can be classified as having "high intrinsic capacity," and that socio-demographic and lifestyle factors such as age, physical activity, tobacco use, and education are strongly linked to IC domains^{20,21}. Meanwhile, analyses of frailty using phenotype models indicate a high prevalence of frailty and pre-frailty ($\approx 29.2\%$ frailty; $\approx 58.8\%$ pre-frailty) among older adults in India, with greater vulnerability observed among women, rural residents, and individuals with lower levels of education.

Despite conceptual agreement and encouraging

early results from high-income and some middle-income regions, there is a notable lack of population-level research from India that explicitly links WHO-defined IC (composite and domain scores) with frailty status. Given India's large demographic scale and within-country diversity, understanding how overall IC and its individual domains relate to frailty in older Indian adults is crucial for guiding integrated, capacity-focused geriatric care models and regionally appropriate public health strategies. Therefore, this study aims to explore the association between intrinsic capacity and frailty among older Indian adults using data from the nationally representative LASI Wave-1.

Methods

LASI main overview

The Longitudinal Ageing Study in India (LASI) utilized a multistage, stratified sampling strategy to ensure both national and state-level representation, drawing upon the 2011 Census as its framework. In this design, districts and tehsils (sub-districts) functioned as primary sampling units, while villages in rural areas and wards in urban areas were selected as secondary units. A detailed description of the study design, including the use of survey weights and sampling methodology, has been previously published¹⁹. For this analysis, we included data from 26,943 community-dwelling individuals aged 60 years and older. All participants gave informed consent prior to participation, having received an information sheet that explained the study objectives, confidentiality procedures, and safety protocols related to health assessments. The first wave of LASI was approved by the Indian Council of Medical Research (ICMR) and adhered to the ethical principles of the Declaration of Helsinki.

Frailty assessment

Frailty was defined using Fried's phenotypic frailty criteria, which include five components: unintentional weight loss, weakness, exhaustion, slowness, and low physical activity. Weight loss was assessed through a self-reported question on unintentional weight loss in the past year. Weakness was evaluated using handgrip strength and classified as low based on sex- and BMI-specific cut-offs. For men, the cut-offs were <29 kg for BMI ≤ 24 , <30 kg for BMI 24.1–26 and 26.1–28, and <32 kg for BMI >28 . For women, the cut-offs were <17 kg for BMI ≤ 23 , <17.3 kg for BMI 23.1–26, <18 kg for BMI 26.1–29, and <21 kg for BMI >29 . Slowness was assessed using a 4-meter gait speed test, with slow gait defined according to sex- and height-specific thresholds. For men, the cut-off was <0.65 m/s for height ≤ 173 cm and <0.76 m/s for height >173 cm. For women, the corresponding cut-offs were <0.65 m/s for height ≤ 159 cm and <0.76 m/s for height >159 cm.

Low physical activity was determined based on a self-reported question regarding the frequency of physical activity, with responses of "never" or "rarely" classified

as low activity. Exhaustion was assessed using the item “everything was an effort,” and participants who responded “often” or “most/all of the time” were considered to have exhaustion. Participants who did not meet any of the five criteria were classified as robust, those meeting one or two criteria were classified as pre-frail, and those meeting three or more criteria were classified as frail.

Intrinsic capacity scoring

IC was assessed across five domains: cognition, locomotor capacity, sensory, vitality, and mood. For the cognition domain, we included multiple components: memory (immediate and delayed recall), orientation (to time and place), arithmetic function (backward counting and computation), executive function (paper folding and pentagon drawing), and object naming. A composite cognitive score was derived by summing the component scores. To account for educational differences, we stratified participants into three groups based on educational attainment: no formal education, education up to middle school, and secondary or higher education. Within each educational category, cognition scores were divided into tertiles, and participants were assigned scores of 0 (lowest tertile), 1 (middle tertile), or 2 (highest tertile), reflecting cognitive capacity adjusted for educational background.

The sensory domain was evaluated based on self-reported difficulty in hearing and vision. Participants with no reported impairment in either domain were scored 2, those with impairment in one domain were scored 1, and those with impairment in both were scored 0. Locomotor capacity was assessed using the semi-tandem balance test. Participants who were able to maintain the position for 10 seconds were assigned a score of 2, those who could maintain it for 3.0 to 9.9 seconds received a score of 1, and those unable to maintain the position for at least 3 seconds were scored 0.

Vitality was assessed using body mass index (BMI). Participants with BMI ≤ 18.4 kg/m² (underweight) were assigned a score of 0, those with BMI between 18.5–24.9 kg/m² (normal weight) were scored 1, and those with BMI ≥ 25.0 kg/m² (overweight or obese) were scored 2. The mood domain was assessed using two questions: “Have you felt sad, blue, or depressed for two weeks or more in a row?” and “Have you lost interest in most things like hobbies, work, or activities?” Participants who responded “yes” to both questions were scored 0, “yes” to one question were scored 1, and “no” to both were scored 2.

The scores from all five domains were summed to generate a total intrinsic capacity score, ranging from 0 to 10, with higher scores indicating better overall intrinsic capacity.

Covariates

Covariates included age, sex (male or female), body mass index (BMI), place of residence (rural or urban), and

living arrangements. Living arrangements were categorized as: living with spouse and children, living with spouse or children, living with others, and living alone. Socioeconomic status was assessed using monthly per capita expenditure (MPCE) quintiles. Marital status and education level were also included, with education categorized as: no formal education, primary to middle school (Class I to VIII), higher secondary or diploma, and graduation or post-graduation. Substance use variables included current use of tobacco and alcohol. Comorbid conditions were based on self-reported physician diagnoses and included diabetes, hypertension, heart disease, stroke, high cholesterol, bone and joint disorders, and chronic lung disease. Multimorbidity was defined as the presence of two or more of these chronic conditions.

Statistical Analysis

Statistical analyses were conducted using Stata version 17.0 (StataCorp. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC; 2021), and graphs were generated using GraphPad Prism version 8 (GraphPad Software, San Diego, CA). Descriptive statistics were used to summarize the study variables. Categorical variables were presented as absolute frequencies and percentages, while continuous variables were reported as mean with standard deviation (SD) or median with interquartile range (IQR), depending on the distribution. The Shapiro–Wilk test was used to assess normality.

Associations between frailty status (categorized as robust, pre-frail, and frail) and categorical variables were examined using the chi-squared test. For continuous variables, one-way analysis of variance (ANOVA) or the Kruskal–Wallis test was used, based on the normality of the data. To explore the association between frailty status and total intrinsic capacity score and individual IC domains, multinomial logistic regression analysis was performed, using the robust group as the reference category. Three models were constructed: Model 1 was unadjusted; Model 2 was adjusted for age and sex; and Model 3 was further adjusted for age, sex, monthly per capita expenditure (MPCE), and the presence of diabetes, hypertension, heart disease, stroke, high cholesterol, bone and joint disease, and chronic lung disease. Self-rated health was not included in adjusted models because it was considered a global health-status indicator with substantial conceptual overlap with both frailty and intrinsic capacity, raising the possibility of overadjustment. Results are reported as relative risk ratios (RRRs) with 95% CIs, as appropriate. A p-value of <0.05 was considered statistically significant.

Results

Baseline characteristics

This analysis included 26,943 community-dwelling older adults (Figure 1). The median age (interquartile




Figure 1. Flow chart depicting the inclusion of participants.




Figure 2. Bar and whisker plot showing intrinsic capacity score between frailty groups.

Table 1. Baseline characteristics of study population (n= 26,943)

Variables	Robust (n=792)	Pre-frail (n= 16,488)	Frail (n=9,663)	p-value
Age [Median (IQR)]	65 (62-68)	65 (62-70)	70 (65-76)	<0.001
60-74 years	743 (93.81)	14382 (87.23)	6585 (68.15)	<0.001
75 years and above	49 (6.19)	2106 (12.77)	3078 (31.85)	
Sex				<0.001
Male	451 (56.94)	8402 (50.96)	4192 (43.42)	
Female	341 (43.06)	8086 (49.04)	5467 (56.58)	
Body mass index [Median (IQR)]	23.58 (21.06-25.81)	21.81 (18.98-24.95)	20.81 (18.07-24.45)	<0.001
Residence				<0.001
Rural	412 (52.02)	10914 (66.19)	6696 (69.30)	
Urban	380 (47.98)	5574 (33.81)	2967 (30.70)	
Living arrangements				<0.001
Living with spouse and children	444 (56.06)	7971 (48.34)	3486 (36.08)	
Living with children or spouse	312 (39.39)	7083 (42.96)	4923 (50.95)	
Living with others	17 (2.15)	691 (4.19)	617 (6.39)	
Living alone	19 (2.40)	743 (4.51)	637 (6.59)	
MPCE				<0.001
Poorest	117 (14.77)	3251 (19.72)	2156 (22.31)	
Poorer	144 (18.18)	3379 (20.49)	2057 (21.29)	
Middle	171 (21.59)	3399 (20.61)	1990 (20.59)	
Richer	158 (19.95)	3312 (20.09)	1848 (19.12)	
Richest	202 (25.51)	3147 (19.09)	1612 (16.68)	
Marital Status				<0.001
Married	633 (79.92)	11564 (70.14)	5349 (55.36)	
Widowed	152 (19.19)	4603 (27.92)	4108 (42.51)	
Divorced	6 (0.76)	184 (1.12)	117 (1.21)	
Never married	1 (0.13)	137 (0.83)	89 (0.92)	
Education				<0.001
No formal education	245 (30.93)	8172 (49.56)	5931 (61.38)	
I to VIII standard	265 (33.46)	5524 (33.50)	2767 (28.63)	
IX-X	108 (13.64)	1401 (8.50)	521 (5.39)	
Higher secondary or diploma	63 (7.95)	644 (3.91)	224 (2.32)	
Graduation or post-graduation	111 (14.02)	747 (4.53)	220 (2.28)	
Tobacco use	274 (34.64)	6639 (40.28)	3763 (38.95)	0.002
Alcohol	148 (18.69)	3019 (18.31)	1524 (15.77)	<0.001
Self-rated health				<0.001
Poor	39 (4.92)	1627 (9.87)	1745 (18.06)	
Fair	210 (26.52)	5157 (31.28)	3478 (36.00)	
Good	320 (40.40)	6456 (39.16)	3153 (32.63)	
Very good	169 (21.34)	2714 (16.46)	1100 (11.38)	
Excellent	54 (6.82)	532 (3.23)	186 (1.93)	

Table 1. (Cont. from previous page).

Variables	Robust (n=792)	Pre-frail (n=16,488)	Frail (n=9,663)	p-value
Comorbidities				
Diabetes	140 (17.68)	2508 (15.22)	1385 (14.33)	0.015
Hypertension	288 (36.36)	11089 (32.73)	6160 (36.24)	<0.001
Heart disease	43 (5.43)	801 (4.86)	485 (5.02)	0.683
Stroke	9 (1.14)	255 (1.55)	291 (3.01)	<0.001
High cholesterol	45 (5.68)	634 (3.85)	315 (3.26)	0.001
Bone and joint disease	115 (14.52)	2588 (15.70)	1968 (20.37)	<0.001
Chronic lung disease	44 (5.56)	1111 (6.74)	870 (9.00)	<0.001
Multimorbidity	181 (22.85)	3565 (21.62)	2393 (24.76)	<0.001

Table 2. The relation between frailty and intrinsic capacity and individual domains.

Variables	Robust (n=792)	Pre-frail (n=16,488)	Frail (n=9,663)	p-value
Total IC score [Median (IQR)]	7 (6-8)	7 (6-8)	6 (5-7)	<0.001
Total IC score [Mean (SD)]	7.08 (1.29)	6.57 (1.43)	6.09 (1.57)	<0.001
Cognition				
Good	391 (49.37)	6729 (40.81)	3031 (31.37)	<0.001
Fair	301 (38.01)	6228 (37.77)	3474 (35.95)	
Poor	100 (12.63)	3531 (21.42)	3158 (32.68)	
Sensory				
Both intact	49 (6.19)	987 (5.99)	844 (8.73)	<0.001
One impaired	459 (57.95)	8287 (50.26)	4896 (50.67)	
Both impaired	284 (35.86)	7214 (43.75)	3923 (40.60)	
Locomotor				
Good	787 (99.37)	16005 (97.07)	8391 (86.84)	<0.001
Fair	5 (0.63)	416 (2.52)	1015 (10.50)	
Poor	0 (0.00)	67 (0.41)	257 (2.66)	
Vitality				
Good	259 (32.70)	4072 (24.70)	2151 (22.26)	<0.001
Fair	458 (57.83)	8995 (54.55)	4729 (48.94)	
Poor	75 (9.47)	3421 (20.75)	2783 (28.80)	
Mood				
Good	676 (85.35)	13683 (83.02)	7365 (76.26)	<0.001
Fair	62 (7.83)	1484 (9.00)	1171 (12.12)	
Poor	54 (6.82)	1315 (7.98)	1122 (11.62)	

range) of the study population was 67 (63-72) years, with nearly half of them male (13,049, 48.43%). Two-thirds of the population (18,022, 66.89%) reside in rural areas. The association between frailty status (robust, pre-frail,

and frail) and demographic factors is reported in Table 1. We observed that a significantly greater proportion of participants aged 75 years and above and females were frail. Frail older adults had significantly lower BMI (20.81

Table 3. Results of multinomial regression analysis Model 3[#].

Variable	Pre-frail		Frail	
	RRR (95% CI)	p-value	RRR (95% CI)	p-value
IC score	0.79 (0.75-0.84)	<0.001	0.67 (0.64-0.71)	<0.001
Individual domains				
Cognition				
Good	1 (ref)		1 (ref)	
Fair	1.14 (0.97-1.33)	0.099	1.27 (1.09-1.49)	0.003
Poor	1.75 (1.39-2.19)	<0.001	2.52 (2.00-3.16)	<0.001
Sensory				
Both intact	1 (ref)		1 (ref)	
One impaired	0.95 (0.69-1.28)	0.727	0.78 (0.57-1.07)	0.122
Both impaired	1.32 (0.96-1.81)	0.086	1.12 (0.81-1.54)	0.499
Locomotor				
Good	1 (ref)		1 (ref)	
Fair	3.29 (1.35-7.98)	0.009	9.79 (4.04-23.74)	<0.001
Poor	-	-	-	-
Vitality				
Good	1 (ref)		1 (ref)	
Fair	1.19 (1.01-1.40)	0.041	1.11 (0.94-1.32)	0.213
Poor	2.56 (1.95-3.36)	<0.001	3.44 (2.61-4.53)	<0.001
Mood				
Good	1 (ref)		1 (ref)	
Fair	1.18 (0.91-1.55)	0.210	1.74 (1.33-2.28)	<0.001
Poor	1.21 (0.91-1.61)	0.177	1.91 (1.44-2.55)	<0.001

Base outcome: Robust. [#]Model 3: adjusted for age, sex, MPCE, diabetes, hypertension, heart disease, stroke, cholesterol, bone and joint disease, chronic lung disease.

in frail vs 21.81 in pre-frail vs 23.58 in robust). Place of residence, living arrangements, socioeconomic status (measured by MPCE), education, and marital status were also associated with frailty. Multimorbidity (presence of two or more comorbidities) was higher among frail older adults ($p < 0.001$).

Intrinsic capacity and frailty

The total IC score was lower among frail older adults [6 (5-7)] compared to pre-frail [7 (6-8)] and robust older adults [7 (6-8)] (Table 2, Figure 2). The analysis of individual IC domains revealed that poor cognition was more common among frail individuals (12.63% vs 21.42% vs 32.68%, $p < 0.001$), while sensory impairment was more prevalent among pre-frail older adults (35.86% vs 43.75% vs 40.60%, $p < 0.001$). Fair locomotor performance (0.63% vs 2.52% vs 10.50%, $p < 0.001$), poor vitality (9.47% vs

20.75% vs 28.80%, $p < 0.001$), and poor mood (6.82% vs 7.98% vs 11.62%, $p < 0.001$) were significantly associated with frailty status. IC score decreased gradually with increasing age (Supplementary Figure 1).

Multinomial regression analysis

The results of multinomial regression analysis (Model 1 and Model 2) are reported in Supplementary Table 1. Model 1 is unadjusted, whereas Model 2 is adjusted for age and sex. Model 3, which is adjusted for age, sex, MPCE, diabetes, hypertension, heart disease, stroke, cholesterol, bone and joint disease, and chronic lung disease, is reported in Table 3. We found that intrinsic capacity (IC score) decreased significantly in pre-frail (RRR: 0.79; 95% CI 0.75-0.84) and frail (RRR: 0.67; 95% CI 0.64-0.71).

Among individual IC domains, poor cognition was associated with an increased risk of being pre-frail (RRR:

1.75; 95% CI 1.39-2.19) and frail (RRR: 2.52; 95% CI 2.00-3.16). Fair locomotor capacity was associated with both pre-frail (RRR: 3.26; 95% CI 1.35-7.98) and frail status (RRR: 9.79; 95% CI 4.04-23.74). Poor vitality was associated with pre-frailty (RRR: 2.56; 95% CI 1.95-3.36) and frailty (RRR: 3.44; 95% CI 2.61-4.53). Poor mood was associated with frailty (RRR: 1.91; 95% CI 1.44-2.55). The sensory domain was not significantly associated with frailty status in our population.

Discussion

In this nationally representative study, we found a strong and consistent association between lower Intrinsic Capacity (IC) and both pre-frailty and frailty. Poor mood and cognition, low vitality, and limited locomotor capacity were the IC domains most significantly associated with frailty. To our knowledge, population-level evidence examining the relationship between intrinsic capacity and frailty in India remains limited. Our study extends the growing international literature by providing evidence from a large nationally representative sample of older Indian adults.

The findings of our current study align with recent literature^{12,16} that demonstrate the close relationship between the two ends of the resilience-vulnerability spectrum in both community and hospital settings. In the study by Shen S et al., declined composite IC scores were associated with frailty [Odds Ratio (OR)=0.45] and a higher risk of falls (OR=0.64) among older hospitalised patients¹². A recent systematic review of 13 observational studies identified adverse IC trajectory patterns associated with increased frailty, mortality, disability, and fall risk²³. In the secondary longitudinal analysis from the 10/66 study across eight LMIC countries, declining IC was linked to a significantly higher risk of frailty, disability, and dementia than in those with high IC, with strong predictive validity (Mortality Hazard Ratio=4.60, 95% CI 4.16; 5.09)¹⁵. Our LASI results extend these findings by offering large-scale, population-level evidence from India.

Poor cognition and impaired mood were independently associated with frailty in our study. This aligns with previous observational studies that have shown psycho-social factors to be closely linked with frailty in older adults^{24,25}. In the longitudinal Multi-domain Alzheimer's Preventive Trial study²⁶, they did not find a significant association between cognitive decline and frailty, but reported a 47% increase in the risk of incident frailty with each additional IC impairment. Conversely, few prospective studies have shown that decline in cognitive capacity is among the earliest predictors of frailty transition in older adults¹⁶. Additionally, with the emergence of the concept of cognitive frailty²⁷, several recent longitudinal studies have reported a significant association between frailty trajectories and incident cognitive decline^{28,29}. These findings emphasise the bi-directional relationship and synergistic effect of frailty and IC impairment on adverse health outcomes in

older adults.

The association between poor vitality and frailty in our study also resonates with previous research highlighting malnutrition and sarcopenia as common biological bases linking IC loss and frailty¹⁸. In the existing literature, vitality or energy homeostasis has been suggested as the highest hierarchical level of the physiological system, which can serve as a biological basis for the other IC domains^{30,31}. Yu et al., in their research, proposed that vulnerability in an older person's vitality can lead to a gradual decline in other IC domains, resulting in different frailty phenotypes and severity¹⁸.

The current study found that a decline in locomotor capacity is associated with both pre-frail and frail status. This agrees with earlier studies¹⁶, which suggest that locomotor impairment can contribute to slowness, a key component of physical frailty. Mobility is considered a core part of the disabling cascade³², and Gonzalez-Bautista E et al., in the 10/66 study, highlighted that mobility impairment can serve as an early clinical marker of this cascade¹⁵. Although some conceptual overlap exists between locomotor performance and frailty phenotype components, the association observed in our study reinforces the importance of mobility as a marker of declining physiological reserve. The locomotor domain demonstrated a pronounced ceiling effect, with nearly all robust participants classified as having good locomotor function and none classified as having poor locomotion. This resulted in sparse data for some locomotor categories and prevented estimation of relative risk ratios for poor locomotor status because of complete separation.

Unlike cognition, locomotion, vitality, and mood, sensory impairment was not independently associated with frailty after multivariable adjustment. This finding suggests that sensory deficits may have a weaker or more indirect relationship with frailty than other intrinsic-capacity domains. While sensory impairment has been reported as a risk factor for functional decline and falls in studies^{33,34}, its association with development of frailty may depend on context, varying with cultural and environmental compensatory mechanisms¹⁴. It is also possible that the self-reported sensory measures available in LASI were insufficiently sensitive to detect clinically meaningful impairment.

The sociodemographic gradients observed in our study are also noteworthy. Frailty was more prevalent among women, the oldest-old, rural residents, and those with lower education and socioeconomic status, consistent with previous Indian studies^{20,22}. Importantly, these groups have been shown to have lower IC levels²¹. This suggests that interventions aimed at maintaining or improving IC could be especially beneficial for disadvantaged subpopulations.

Our nationwide population-based current study provides valuable insight into the intricate relationship between IC and frailty in community-dwelling older adults in India. While

future waves of LASI are needed to shed light on the causal link between these trajectories, the study also emphasises the utility of IC as a positive, capacity-oriented framework to complement frailty screening and interventions in geriatric practice. The VIVIFRAIL multicomponent exercise program has been found to be an effective strategy to strengthen IC, especially in the domains of locomotion, vitality, and cognition³⁵. Incorporating IC assessments into India's National Programme for the Healthcare of the Elderly (NPHCE) could support proactive interventions to delay frailty progression, extend functional ability, and reduce the healthcare burden, but longitudinal studies are needed to establish temporal relationships and evaluate predictive validity before widespread implementation.

Strengths and Limitations

This study has several strengths. First, it uses data from the LASI, which employed a multistage, stratified sampling design to ensure national and state-level representativeness of older adults, thereby enhancing the generalizability of the findings. Second, frailty was assessed using Fried's phenotypic criteria, a well-established and validated framework that allows for comparability with international studies. Third, IC was comprehensively evaluated across all five domains—cognition, locomotor capacity, sensory, vitality, and mood—using structured assessments, and a total IC score was generated to capture overall functional reserve. Additionally, cognitive scoring was adjusted for educational attainment by categorizing participants into tertiles within education strata, helping to minimize the confounding effect of educational disparities on cognitive assessment. The use of multinomial logistic regression with sequentially adjusted models and inclusion of a broad set of sociodemographic and clinical covariates further strengthens the analytical robustness of the study.

However, several limitations should be noted. The cross-sectional nature of the data precludes causal inference and limits the ability to assess changes in frailty or IC over time. Many key variables, including comorbidities, physical activity, and sensory impairments, were based on self-report and may be prone to recall bias or misclassification. Some overlap exists between components of Fried's frailty phenotype and specific intrinsic-capacity domains, particularly locomotion and vitality. Consequently, part of the observed association may reflect shared underlying constructs. However, intrinsic capacity additionally incorporates cognition, sensory function, and psychological health, which are not included in Fried's phenotype. Therefore, the observed associations likely reflect both shared and distinct dimensions of healthy ageing. Some IC domains, such as mood and sensory function, were assessed using limited questions rather than comprehensive clinical tools, which may affect the precision of domain-specific scoring. Although the models adjusted for a wide range of covariates, residual confounding from unmeasured factors

such as nutritional status, inflammation, or sarcopenia cannot be ruled out. Finally, the absence of longitudinal follow-up data limits the ability to evaluate trajectories of frailty and intrinsic capacity over time; future waves of LASI will be essential to address these gaps.

Conclusion

Intrinsic capacity is strongly and inversely associated with frailty among older Indian adults. Declines in cognition, locomotion, vitality, and mood are key markers of frailty. These findings highlight the potential of incorporating IC-based assessments into geriatric care to identify high-risk individuals and guide early interventions. Capacity-oriented approaches may play a crucial role in promoting healthy ageing and delaying frailty in India's older population.

Ethics approval

The data is freely available in the public domain and survey agencies that conducted the field survey for the data collection have collected prior consent from the respondents. The Indian Council of Medical Research (ICMR) and all partner institutions extended the necessary guidance and ethical approval for conducting the LASI. The LASI data can be accessed for free at the International Institute for Population Sciences website (<https://www.iipsindia.ac.in/content/LASI-data>).

Consent to participate

The survey agencies that conducted the field survey for the data collection have collected prior informed consent from all subjects and/or their legal guardian(s).

Authors' contributions

All authors met the criteria for authorship as follows: study concept and design (Sudeep M. George, Kritartha Kashyap), acquisition of data (Abhijith R. Rao), analysis and interpretation of data (Abhijith R. Rao), and preparation of manuscript (Sudeep M. George, Kritartha Kashyap, Abhijith R. Rao, Minakshi Dhar, Vasu Digra, Parul Bhutani, Pankhuri Saxena, Rajkumar Tata, Bhabhor K Narsinhbhai, Shreya Biswal, Pramod K Mehta, Prasun Chatterjee, Avinash Chakrawarty, Naveet Wig.). No unnamed contributor played a role in manuscript preparation. All the authors read and approved the final version of the manuscript and accept responsibility for the integrity and accuracy of the article's content.

Funding

The LASI project is jointly funded by the National Institute on Aging (NIA) of the National Institutes of Health (NIH), the Government of India (GoI), and the United Nations Population Fund (UNFPA), India. However, the authors received no specific funding for this study.

Acknowledgements

All the authors thank the original data collectors,

depositors, copyright holders, and funders of the Longitudinal Ageing Study in India (LASI).

The editing of this paper was supported by generative AI tools (ChatGPT and Grammarly). ChatGPT was used on selected sections of the manuscript to reduce redundancy to fit into the journal's word limit and enhance fluency without altering the content. In doing so, we uploaded the selected sections identified as long and with unnecessary content and asked ChatGPT to cut the redundancies and enhance the fluency. Grammarly (embedded in Microsoft Word) was used to ensure grammar, spelling and phrasing accuracy. The whole process was human-supervised, and all content has been carefully inspected and revised by authors to confirm accuracy.

References


- United Nations, Department of Economic and Social Affairs, Population Division. World Population Ageing 2020 Highlights: Living arrangements of older persons. New York: United Nations; 2020.
- López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013 Jun 6;153(6):1194-217.
- Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*. 2013 Mar 2;381(9868):752-62. Epub 2013 Feb 8. Erratum in: *Lancet*. 2013 Oct 19;382(9901):1328.
- Fried LP, Tangen CM, Walston J, et al; Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001 Mar;56(3):M146-56.
- World Health Organization. World report on ageing and health. Geneva: WHO; 2015. (Defines intrinsic capacity and introduces the healthy-ageing framework). Available from: <https://apps.who.int/iris/handle/10665/186468>.
- Beard JR, Officer A, de Carvalho IA, et al. The World report on ageing and health: a policy framework for healthy ageing. *Lancet*. 2016 May 21;387(10033):2145-2154.
- Seals DR, Justice JN, LaRocca TJ. Physiological geroscience: targeting function to increase healthspan and achieve optimal longevity. *J Physiol*. 2016 Apr 15;594(8):2001-24.
- Sánchez-Sánchez JL, et al. Association of intrinsic capacity with functional decline and adverse outcomes: a systematic review and meta-analysis. *Lancet Healthy Longevity*. 2024
- Belloni G, Cesari M. Frailty and Intrinsic Capacity: Two Distinct but Related Constructs. *Front Med (Lausanne)*. 2019 Jun 18;6:133.
- Cesari M, Canevelli M, Amuthavalli Thiyagarajan J, Arai H, Assantachai P, Chan P, et al. Frailty and intrinsic capacity: integrating complementary concepts to promote healthy ageing and transformation of care. *Age Ageing*. 2026;55:afag138.
- Stefano Cacciatore, Intrinsic capacity will not replace frailty—and should not try, *Age and Ageing*, Volume 55, Issue 5, May 2026, afag137
- Shen S, Xie Y, Zeng X, et al. Associations of intrinsic capacity, fall risk and frailty in old inpatients. *Front Public Health*. 2023 Oct 10;11:1177812.
- Tay L, Tay EL, Mah SM, Latib A, Koh C, Ng YS. Association of Intrinsic Capacity with Frailty, Physical Fitness and Adverse Health Outcomes in Community-Dwelling Older Adults. *J Frailty Aging*. 2023;12(1):7-15.
- Zhou J, Chang H, Leng M, Wang Z. Intrinsic Capacity to Predict Future Adverse Health Outcomes in Older Adults: A Scoping Review. *Healthcare (Basel)*. 2023 Feb 4;11(4):450.
- Gonzalez-Bautista E, Llibre-Guerra JJ, Sosa AL, et al. Exploring the natural history of intrinsic capacity impairments: longitudinal patterns in the 10/66 study. *Age Ageing*. 2023 Jul 1;52(7):afad137.
- Jia, S., Zhao, W., Ge, M. et al. Associations between transitions of intrinsic capacity and frailty status, and 3-year disability. *BMC Geriatr* 23, 96 (2023).
- Cacciatore S, Marzetti E, Calvani R, et al. Intrinsic capacity and recent falls in adults 80 years and older living in the community: results from the iLSIRENTE Study. *Aging Clin Exp Res*. 2024 Aug 10;36(1):169.
- Yu Z, Lozupone M, Chen J, Bao Z, Ruan Q, Panza F. The Biological Rationale for Integrating Intrinsic Capacity into Frailty Models. *Clin Interv Aging*. 2025 Mar 4;20:273-286.
- International Institute for Population Sciences (IIPS), National Programme for Health Care of Elderly (NPHCE), MoHFW, Harvard T. H. Chan School of Public Health (HSPH) and the University of Southern California (USC) 2020. Longitudinal Ageing Study in India (LASI) Wave 1, 2017-18, India Report, International Institute for Population Sciences, Mumbai.
- Muneera K, Muhammad T, Pai M, Ahmed W, Althaf S. Associations between intrinsic capacity, functional difficulty, and fall outcomes among older adults in India. *Sci Rep*. 2023 Jun 17;13(1):9829.
- Muneera, K., Muhammad, T. & Althaf, S. Socio-demographic and lifestyle factors associated with intrinsic capacity among older adults: evidence from India. *BMC Geriatr* 22, 851 (2022).
- Nagarkar A, Kulkarni AS. Regional variation in prevalence of frailty in India: Evidence from longitudinal ageing study in India (LASI) wave-1. *Indian J Med Res*. 2024 May;159(5):441-448.
- Wang Y, Tang N, Shao M, Song J, Su Q, Gao Y. Transitions and trajectories in intrinsic capacity states over time: a systematic review. *BMC Geriatr*. 2025 Jun 4;25(1):407.
- Brennan, T.H., Lewis, L.K., Gordon, S.J. et al. Behavioural and psychological factors associated with pre-frailty in community-dwelling adults aged 40 and over: a cross-sectional study. *BMC Public Health* 25, 956 (2025).
- Żółkowska B, Lee CS, Denfeld QE, et al. Clinical and Psychological Factors Associated with Frailty in Patients with Heart Failure. *J Clin Med*. 2024 Dec 2;13(23):7345.
- Gonzalez-Bautista E, de Souto BP, Andrieu S, Rolland Y, Vellas B, group MD. Screening for intrinsic capacity impairments as markers of increased risk of frailty and disability in the context of integrated care for older people: secondary analysis of MAPT. *Maturitas*. (2021) 150:1-6.
- Kelaiditi E, Cesari M, Canevelli M, et al. Cognitive frailty: rational and definition from an (I.A.N.A./I.A.G.G.) international consensus group. *J Nutr Health Aging*. 2013;17(9):726-34.
- Boyle PA, Buchman AS, Wilson RS, Leurgans SE, Bennett DA. Physical frailty is associated with incident mild cognitive impairment in community-based older persons. *J Am Geriatr Soc*. 2010 Feb;58(2):248-55.
- Buchman AS, Boyle PA, Wilson RS, Tang Y, Bennett DA. Frailty is associated with incident Alzheimer's disease and cognitive decline in the elderly. *Psychosom Med*. 2007 Jun;69(5):483-9.
- Beard JR, Jotheeswaran AT, Cesari M, Araujo de Carvalho I. The structure and predictive value of intrinsic capacity in a longitudinal study of ageing. *BMJ open*. 2019;9(11):e026119.
- Cesari M, Araujo de Carvalho I, Amuthavalli Thiyagarajan J, et al. Evidence for the domains supporting the construct of intrinsic capacity. *J Gerontol a Biol Sci Med Sci*. 2018;73(12):1653-

- 1660.
32. Verbrugge LM, Jette AM. The disablement process. Soc Sci Med. 1994 Jan;38(1):1-14.
 33. Tan BKJ, Man REK, Gan ATL, et al. Is Sensory Loss an Understudied Risk Factor for Frailty? A Systematic Review and Meta-analysis. J Gerontol A Biol Sci Med Sci. 2020 Nov 13;75(12):2461-2470.
 34. Yamada Y, Nakashima H, Nagae M, et al. Dual sensory impairment predicts an increased risk of postdischarge falls in older patients. J Am Med Dir Assoc. 2024 Sep;25(9):105123.
 35. Sanchez-Sanchez JL, de Souto BP, Anton-Rodrigo I, et al. Effects of a 12-week Vivifrail exercise program on intrinsic capacity among frail cognitively impaired community-dwelling older adults: secondary analysis of a multicentre randomised clinical trial. Age Ageing. (2022) 51: afac303.

Supplementary Table 1. Results of multinomial regression analysis Model 1 and Model 2.

Variables	Model 1 [#]		Model 2 ^{##}	
	Pre-frail RRR (95% CI)	Frail RRR (95% CI)	Pre-frail RRR (95% CI)	Frail RRR (95% CI)
IC score	0.77 (0.74-0.82)***	0.62 (0.59-0.66)***	0.78 (0.74-0.82)***	0.66 (0.63-0.70)***
Individual domains	RRR (95% CI)	RRR (95% CI)	RRR (95% CI)	RRR (95% CI)
Cognition				
Good	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Fair	1.20 (1.03-1.40)*	1.49 (1.27-1.74)***	1.16 (0.99-1.35)	1.30 (1.11-1.53)**
Poor	2.05 (1.64-2.56)***	4.07 (3.25-3.10)***	1.82 (1.45-2.28)***	2.67 (2.12-3.35)***
Sensory				
Both intact	1 (ref)	1 (ref)	1 (ref)	1 (ref)
One impaired	0.89 (0.66-1.21)	0.62 (0.45-0.84)**	0.96 (0.70-1.30)	0.78 (0.57-1.06)
Both impaired	1.26 (0.92-1.72)	0.80 (0.58-1.09)	0.78 (0.57-1.06)	1.18 (0.86-1.62)
Locomotor				
Good	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Fair	4.09 (1.68-9.91)**	19.04 (7.88-45.99)***	3.27 (1.35-7.94)**	9.94 (4.10-24.07)***
Poor	-	-	-	-
Vitality				
Good	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Fair	1.25 (1.07-1.46)**	1.24 (1.06-1.46)**	1.25 (1.06-1.46)**	1.15 (0.97-1.35)
Poor	2.90 (2.23-3.77)	4.47 (3.43 to 5.81)***	2.82 (2.16-3.66)***	3.70 (2.84-4.83)***
Mood				
Good	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Fair	1.18 (0.91-1.54)	1.73 (1.32-2.26)***	1.18 (0.91-1.54)	1.77 (1.35-2.32)***
Poor	1.20 (0.90-1.59)	1.91 (1.43-2.53)***	1.21 (0.91-1.60)	1.95 (1.47-2.61)***

Base outcome: Robust. [#]Model 1: unadjusted. ^{##}Model 2: adjusted for age and sex. *p-value<0.05. **p-value<0.01. ***p-value<0.001.



Supplementary Figure 1. Scatter plot with fitter line showing association between age and intrinsic capacity score.